

# Unconditional Prenatal Income Supplement and Birth Outcomes

Marni D. Brownell, PhD,<sup>a,b</sup> Mariette J. Chartier, PhD,<sup>a,b</sup> Nathan C. Nickel, PhD,<sup>a,b</sup> Dan Chateau, PhD,<sup>a,b</sup> Patricia J. Martens, PhD,<sup>a,b,†</sup> Joykrishna Sarkar, MSc,<sup>b</sup> Elaine Burland, PhD,<sup>b</sup> Douglas P. Jutte, MD, MPH,<sup>c</sup> Carole Taylor, MSc,<sup>b</sup> Robert G. Santos, PhD,<sup>a,b</sup> Alan Katz, MB, ChB,<sup>a,b</sup> On behalf of the PATHS Equity for Children Team

## abstract

**BACKGROUND AND OBJECTIVES:** Perinatal outcomes have improved in developed countries but remain poor for disadvantaged populations. We examined whether an unconditional income supplement to low-income pregnant women was associated with improved birth outcomes.

**METHODS:** This study included all mother–newborn pairs (2003–2010) in Manitoba, Canada, where the mother received prenatal social assistance, the infant was born in the hospital, and the pair had a risk screen ( $N = 14\,591$ ). Low-income women who received the income supplement (Healthy Baby Prenatal Benefit [HBPB],  $n = 10\,738$ ) were compared with low-income women who did not receive HBPB ( $n = 3853$ ) on the following factors: low birth weight, preterm, small and large for gestational age, Apgar score, breastfeeding initiation, neonatal readmission, and newborn hospital length of stay (LOS). Covariates from risk screens were used to develop propensity scores and to balance differences between groups in regression models;  $\gamma$  sensitivity analyses were conducted to assess sensitivity to unmeasured confounding. Population-attributable and preventable fractions were calculated.

**RESULTS:** HBPB was associated with reductions in low birth weight (aRR, 0.71 [95% CI, 0.63–0.81]), preterm births (aRR, 0.76 [95% CI, 0.69–0.84]) and small for gestational age births (aRR, 0.90 [95% CI, 0.81–0.99]) and increases in breastfeeding (aRR, 1.06 [95% CI, 1.03–1.09]) and large for gestational age births (aRR, 1.13 [95% CI, 1.05–1.23]). For vaginal births, HBPB was associated with shortened LOS (weighted mean, 2.86;  $P < .0001$ ). Results for breastfeeding, low birth weight, preterm birth, and LOS were robust to unmeasured confounding. Reductions of 21% (95% CI, 13.6–28.3) for low birth weight births and 17.5% (95% CI, 11.2–23.8) for preterm births were associated with HBPB.

**CONCLUSIONS:** Receipt of an unconditional prenatal income supplement was associated with positive outcomes. Placing conditions on income supplements may not be necessary to promote prenatal and perinatal health.

<sup>a</sup>Department of Community Health Sciences, University of Manitoba, Winnipeg, Manitoba, Canada; <sup>b</sup>Manitoba Centre for Health Policy, Winnipeg, Manitoba, Canada; and <sup>c</sup>School of Public Health, University of California, Berkeley, Berkeley, California

<sup>†</sup>Deceased.

Dr Brownell conceptualized and designed the study, interpreted the data, and drafted the manuscript; Drs Chartier, Nickel, Chateau, Burland, Jutte, Santos, and Katz contributed to the conceptualization and design of the study, as well as the interpretation of the data, and critically reviewed drafts of the manuscript; Dr Martens contributed to the conceptualization and design of the study, as well as the interpretation of the data, and critically reviewed drafts of the manuscript; she did not approve the final manuscript as submitted because she died in 2015;

**WHAT'S KNOWN ON THIS SUBJECT:** Perinatal outcomes have improved in developed countries but remain poor for socioeconomically disadvantaged populations. Evaluations of conditional income supplement programs for low-income pregnant women have yielded mixed results because of methodologic challenges.

**WHAT THIS STUDY ADDS:** Using propensity scores to balance exposed and unexposed groups, we found that an unconditional prenatal income supplement was associated with positive perinatal results. Placing conditions on income supplements may not be necessary to promote prenatal and perinatal health.

**To cite:** Brownell MD, Chartier MJ, Nickel NC, et al. Unconditional Prenatal Income Supplement and Birth Outcomes. *Pediatrics*. 2016;137(6):e20152992

The prenatal period is crucial in terms of both newborn and lifelong health.<sup>1-3</sup> Prenatal exposure to factors (including severe stress, poor nutrition, and substance use) can lead to adverse birth outcomes such as low birth weight and preterm births, which have an impact on health and development throughout childhood and beyond.<sup>2,4-17</sup> Women living in poverty are more likely to be exposed to high levels of stress, have inadequate nutritional intake, and smoke and/or drink or use substances during pregnancy; they are also more likely to give birth to preterm or low birth weight infants.<sup>18</sup>

Considerable focus has been placed on improving outcomes for infants born to women living in poverty, through the use of prenatal interventions, in both developed and developing countries. There are several programs in Latin America, including Oportunidades in Mexico<sup>19</sup> and the Bolsa Familia Program in Brazil,<sup>20</sup> that provide money conditional on certain behaviors such as attending prenatal care.<sup>19-21</sup> Although many of these programs are not administered specifically during the prenatal period, they have been found to influence birth outcomes.<sup>22</sup> A Cochrane Review of 10 evaluations of conditional cash transfer programs concluded that although there was strong evidence of a positive impact on health service utilization and health outcomes, it was difficult to determine the role the cash incentives played in the outcomes.<sup>21</sup> An examination of participants in the Oportunidades program concluded that it was the cash itself that was leading to health benefits.<sup>19</sup>

In the United States, programs aimed at promoting prenatal health for women living in poverty have followed a different model, typically offering free services rather than conditional cash transfers. One of the best known, the Special Supplemental Nutrition Program for Women, Infants, and Children

(WIC), targets low-income women in the prenatal and postnatal periods and provides food supplementation, nutrition education, and access to health care services.<sup>23</sup> Evaluations of WIC have yielded mixed results, mainly due to challenges with identifying comparable exposed and unexposed groups.<sup>24-29</sup> An evaluation using propensity score matching concluded that previous positive associations between birth outcomes and WIC may have been exaggerated due to estimation methods that did not account for unmeasured confounding.<sup>27</sup>

In 2001, the Canadian province of Manitoba introduced the Healthy Baby Prenatal Benefit (HBPB) to improve prenatal health and birth outcomes. Within Canada's universal health care, prenatal care is already provided free of cost. HBPB provides prenatal income support (up to Can\$81.41 monthly) to low-income women during the second and third trimesters. HBPB is unique in that the income supplement is provided without any conditions. Although pamphlets about the importance of good prenatal nutrition and information about breastfeeding and healthy infant development accompany the mailed monthly payment, women can spend the money as they see fit. The objective of the present study was to determine whether an unconditional income supplement to low-income pregnant women was associated with improved birth outcomes.

## METHODS

### Population and Data Source

This study was conducted at the Manitoba Centre for Health Policy, as part of the PATHS Equity for Children program of research,<sup>30</sup> and received approval from the University of Manitoba's Health Research Ethics Board. Data came from the PATHS Data Resource, which collects population-wide, de-identified

health and social services data for children registered for the universal health care program in Manitoba (population, 1.2 million).<sup>30-33</sup> The databases used in this study included HBPB program data, newborn risk screen data, hospital discharge abstracts, social assistance (ie, welfare), physician visit records, prescription medication records, a population health registry, and the Canadian census.

All low-income pregnant women are eligible to apply for HBPB, and those applicants with documented annual incomes below Can\$32 000 whose pregnancy has been confirmed by a physician are enrolled. The Healthy Child Manitoba Office, which administers HBPB, maintains administrative data on all applicants and recipients. They also maintain a database of information about families with newborns from a universal risk screen that is administered shortly after birth by public health nurses.<sup>34</sup> This screen provides information about prenatal health and health behaviors as well as social risk factors. Both these databases are linkable at the individual level to the population-wide information on health and social service use, using a scrambled identifier. The ability to combine program participation information with information on family risk factors and service use presented an exceptional opportunity to evaluate the impact of HBPB by enabling us to ensure comparability between those exposed and not exposed. The validity of data in the PATHS Data Resource has been well documented.<sup>32,35-40</sup>

The initial study population included all mother-infant pairs for Manitoba women who had a live hospital birth from January 1, 2003, through December 31, 2010 (Fig 1); <1% of Manitoba births occur at home.<sup>41</sup> A quasi-experimental retrospective cohort design was used, comparing birth outcomes for infants of women

who received (exposed) or did not receive (unexposed) HBPB. Although almost one-third of pregnant women are eligible for HBPB,<sup>42</sup> we selected all women receiving welfare during pregnancy ( $N = 16\,557$ ) to identify comparable exposed and unexposed groups; almost one-half of those eligible for HBPB receive welfare. Pregnant women receiving welfare represent a very-low-income population, requiring help to meet basic personal and family needs, and are therefore a group at risk for poor birth outcomes. Preliminary analyses found that the exposed and unexposed groups of women receiving welfare during pregnancy had comparable low mean annual incomes at Can\$9941 and Can\$9972, respectively; HBPB represents an almost 10% increase in their monthly income. Although all women receiving welfare during pregnancy are eligible for HBPB, not all apply. Reasons for not applying are not recorded but could affect the comparability between the exposed and unexposed groups. The newborn risk screen was developed and validated for predicting families at risk for maltreating their children.<sup>34</sup> Screen data are available for almost all families with newborns in the province, and they contain detailed information on factors such as health behaviors (eg, prenatal smoking, alcohol consumption), maternal mental health, and family functioning. Information from newborn risk screens was used to ensure comparability of the exposed and unexposed groups.

## Measures

The exposure variable was whether the mother received HBPB. Because the preliminary analyses revealed that almost all HBPB recipients in our study (99.1%) received the maximum benefit, a dose-response effect was not examined. Information on birth outcomes was extracted from hospital discharge abstracts.


**TABLE 1** Outcomes Examined for Exposed (Received HBPB) and Unexposed (Did Not Receive HBPB) Groups

| Outcome                    | Definition                                                                                                                                                        |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Low birth weight           | <2500 g                                                                                                                                                           |
| Preterm birth              | <37 wk gestational age                                                                                                                                            |
| Small for gestational age  | <10th percentile for gestational age and gender, using a Canadian standard <sup>43</sup>                                                                          |
| Large for gestational age  | >90th percentile for gestational age and gender, using a Canadian standard <sup>43</sup>                                                                          |
| 5-min Apgar score          | Dichotomized into $\leq 7$ and $\geq 8$                                                                                                                           |
| Breastfeeding initiation   | Exclusive or partial breastfeeding at hospital discharge                                                                                                          |
| Neonatal readmission       | Readmission to any Manitoba hospital within 28 d of birth                                                                                                         |
| LOS, birth hospitalization | LOS at birth; continuous measures calculated separately for infants delivered vaginally and by cesarean delivery due to different stays expected for these groups |

Definitions of outcomes examined are given in Table 1.

A number of covariates were available for the exposed and unexposed groups (Table 2). Most of

these were taken from the newborn risk screen and were answered yes or no; where information was missing, a third category ("missing") was added. Additional confounders analyzed were: maternal diabetes, defined



**FIGURE 1** Selection of low-income groups exposed and not exposed to the HBPB.

through a combination of hospital visit, physician visit, and medication records<sup>44,45</sup>; mother's age at birth of first child,<sup>46</sup> from the population health registry; and an index of area-level socioeconomic status<sup>47</sup> compiled from Canada census data.

### Statistical Analysis

Due to the potential for systematic differences between women on welfare who did or did not apply for HBPB, propensity scores were used to adjust for measured confounding.<sup>48,49</sup> A woman's propensity score is her probability of receiving HBPB, given her measurable characteristics. Adjusting for the propensity score is an efficient strategy to test for and balance observed differences between those receiving and not receiving HBPB. Propensity scores allow one to make comparisons between similar exposed and unexposed groups. Propensity scores were estimated by using multiple logistic regression, with HBPB as the dependent variable and the covariates presented in Table 2. The estimated propensity scores were used to construct inverse probability of treatment weights (IPTWs). IPTWs were applied to the data to balance differences in observed characteristics between HBPB recipients and nonrecipients. We tested whether the measured confounding covariates were balanced by using standardized differences,<sup>50–52</sup> set at an a priori 10% difference.<sup>50</sup> Once we achieved balance in measured covariates, IPTWs were applied to all outcome models to estimate the adjusted association between receipt of HBPB and the outcomes.

### Outcome Models

Dichotomous outcomes were modeled by using generalized linear models with a binomial distribution. The log-link function was used to estimate the risk ratio associated with receiving HBPB for each outcome. We first modeled

**TABLE 2** Characteristics of Women on Welfare During Pregnancy According to Receipt of HBPB

| Characteristic                                | Before IPTW Applied |                      |          |                          | After IPTW Applied: Standardized Differences |
|-----------------------------------------------|---------------------|----------------------|----------|--------------------------|----------------------------------------------|
|                                               | Received HBPB       | Did Not Receive HBPB | <i>P</i> | Standardized Differences |                                              |
| Categorical variables                         |                     |                      |          |                          |                                              |
| Received screen prenatally <sup>a</sup>       | 1074 (10.0)         | 192 (5.0)            | <.001    | 19.15                    | 0.14                                         |
| Alcohol use during pregnancy                  | 2378 (22.2)         | 789 (20.5)           | .03      | 4.07                     | 1.07                                         |
| Drug use during pregnancy                     | 1446 (13.5)         | 456 (11.8)           | .01      | 4.91                     | 0.37                                         |
| Family history of disability                  | 375 (3.5)           | 136 (3.5)            | .91      | 0.20                     | 0.69                                         |
| Smoked during pregnancy                       | 4954 (46.1)         | 1783 (46.3)          | .88      | 0.28                     | 0.11                                         |
| Mother did not complete high school           | 4932 (45.9)         | 1628 (42.3)          | <.001    | 7.41                     | 0.61                                         |
| Received welfare                              | 6863 (63.9)         | 2001 (51.9)          | <.001    | 24.45                    | 0.41                                         |
| Single-parent family                          | 4722 (44.0)         | 1376 (35.7)          | <.001    | 16.94                    | 0.75                                         |
| No prenatal care before 6 mo                  | 626 (5.8)           | 430 (11.2)           | <.001    | 19.21                    | 0.14                                         |
| Mother has depression                         | 2110 (19.7)         | 602 (15.6)           | <.001    | 10.58                    | 0.31                                         |
| Mother has anxiety disorder                   | 719 (6.7)           | 196 (5.1)            | <.001    | 6.84                     | 1.17                                         |
| Mother has schizophrenia                      | 130 (1.2)           | 38 (1.0)             | .24      | 2.15                     | 0.36                                         |
| Mother has a mental disability                | 136 (1.3)           | 45 (1.2)             | .63      | 0.90                     | 0.51                                         |
| Antisocial father                             | 402 (3.7)           | 95 (2.5)             | <.001    | 7.37                     | 0.93                                         |
| Antisocial mother                             | 182 (1.7)           | 55 (1.4)             | .24      | 2.16                     | 0.27                                         |
| Current substance abuse by mother             | 283 (2.6)           | 100 (2.6)            | .89      | 0.25                     | 0.22                                         |
| Social isolation                              | 692 (6.4)           | 248 (6.4)            | .99      | 0.03                     | 0.19                                         |
| Relationship distress                         | 1631 (15.2)         | 472 (12.3)           | <.001    | 8.55                     | 0.46                                         |
| Violence between parents                      | 769 (7.2)           | 211 (5.5)            | <.001    | 6.93                     | 0.27                                         |
| Mother abused as a child                      | 1686 (15.7)         | 504 (13.1)           | <.001    | 7.47                     | 0.52                                         |
| Maternal diabetes                             | 268 (2.5)           | 62 (1.6)             | <.001    | 6.26                     | 0.80                                         |
| Continuous variables                          |                     |                      |          |                          |                                              |
| Mother's age at first birth                   |                     |                      |          | 4.21                     | 1.00                                         |
| Count                                         | 10 738              | 3853                 |          |                          |                                              |
| Mean                                          | 19.8                | 19.6                 |          |                          |                                              |
| Median                                        | 19.1                | 19.0                 |          |                          |                                              |
| SD                                            | 3.4                 | 3.5                  |          |                          |                                              |
| Minimum                                       | 12.0                | 12.2                 |          |                          |                                              |
| Maximum                                       | 42.7                | 43.7                 |          |                          |                                              |
| Area-level SES index (higher value = low SES) |                     |                      |          | 13.40                    | 0.70                                         |
| Count                                         | 10 729              | 3849                 |          |                          |                                              |
| Mean                                          | 0.9                 | 0.8                  |          |                          |                                              |
| Median                                        | 0.8                 | 0.7                  |          |                          |                                              |
| SD                                            | 0.9                 | 0.9                  |          |                          |                                              |
| Minimum                                       | −2.9                | −2.9                 |          |                          |                                              |
| Maximum                                       | 3.9                 | 3.9                  |          |                          |                                              |

Data are presented as *n* (%) or %. SES, socioeconomic status.

<sup>a</sup> Screens are generally completed postnatally, and prenatal screens are indicative of higher risk families.

crude risk ratios and then modeled propensity score–adjusted risk ratios by applying the IPTWs to the dichotomous outcome models.

The 2 continuous hospital length of stay (LOS) outcomes were modeled by using generalized linear models with a negative binomial distribution. The log-link function was used to estimate the ratio in mean LOS associated with exposure to HBPB, first modeling crude mean LOS ratios and then modeling propensity

score–adjusted LOS ratios by applying the IPTWs.

### Sensitivity Analysis

Multiple regression and propensity score methods rest on the assumption that adjustment controls for measured and unmeasured confounding. Although this assumption cannot be directly tested, sensitivity to unmeasured confounding can be assessed.<sup>53</sup> We conducted a  $\gamma$  sensitivity analysis

**TABLE 3** Crude Rates or Means and Relative Risks or Means for Birth Outcomes and Neonatal Readmission for Exposed (Received HBPB) and Unexposed (Did Not Receive HBPB) Groups

| Outcome                      | Crude Rates, %   |         | Risk Ratio             | 95% CI           | P      | Sensitivity to Unmeasured Confounding <sup>a</sup> |
|------------------------------|------------------|---------|------------------------|------------------|--------|----------------------------------------------------|
|                              | HBPB             | No HBPB |                        |                  |        |                                                    |
| Breastfeeding initiation     | 64.2             | 58.9    | 1.06                   | 1.03–1.09        | <.0001 | 56.4                                               |
| Low birth weight (<2500 g)   | 5.1              | 7.8     | 0.71                   | 0.63–0.81        | <.0001 | 61.8                                               |
| Preterm (<37 wk gestation)   | 8.2              | 11.3    | 0.76                   | 0.69–0.84        | <.0001 | 62.9                                               |
| Small for gestational age    | 8.3              | 9.6     | 0.90                   | 0.81–0.99        | .05    | 1.8                                                |
| Large for gestational age    | 16.1             | 13.9    | 1.13                   | 1.05–1.23        | .001   | 38.8                                               |
| Low 5-min Apgar score        | 3.7              | 4.0     | 0.93                   | 0.79–1.09        | .36    | NS                                                 |
| Neonatal readmission (<29 d) | 2.7              | 2.7     | 1.02                   | 0.84–1.25        | .82    | NS                                                 |
|                              | Unweighted Means |         | Weighted Mean (95% CI) |                  |        |                                                    |
| Birth hospital LOS           | HBPB             | No HBPB | HBPB                   | No HBPB          |        |                                                    |
| Births by cesarean delivery  | 7.1              | 7.6     | 7.13 (6.78–7.48)       | 7.17 (6.79–7.55) | .87    |                                                    |
| Vaginal births               | 2.9              | 3.1     | 2.86 (2.79–2.92)       | 3.11 (3.03–3.20) | <.0001 |                                                    |

<sup>a</sup> Analyzed by using  $\gamma$  sensitivity test;  $\gamma$  sensitivity analysis was not calculated for those findings that were not statistically significant (NS).

to answer the question: how strong would any unmeasured confounder have to be to nullify our statistically significant results? Examples of potential unmeasured confounders that might differ between our groups and be associated with newborn outcomes include whether the pregnancy was planned and self-care factors (eg, nutritional intake, stress reduction).

#### Population-Attributable and Population-Preventable Fractions

To quantify the impact of HBPB, population-attributable fractions (PAFs) and population-preventable fractions (PPFs) were calculated. For outcomes in which HBPB was associated with an increase, the PAF was calculated by using the formula  $PAF = Pe \times [(RR - 1)/RR]$ , where  $Pe$  is prevalence of the exposure.<sup>54</sup> For outcomes in which HBPB was associated with a reduction, the PPF was calculated by using the formula  $PPF = Pe \times (1 - RR)$ . For both measures, confidence intervals (CIs) were calculated by using the SD of a bootstrapped mean PAF (or PPF) derived from 500 samples of the population.

## RESULTS

There were 14 591 women who gave birth to live singletons in Manitoba between 2003 and 2010 who had


received welfare during pregnancy and had risk screen information; of these, 10 738 received HBPB, and 3853 did not (Fig 1). Table 2 displays the number and percentage of women in each group for each covariate, as well as the standardized differences between the groups exposed and unexposed to HBPB, before and after applying the IPTWs. Before applying the weights, the standardized differences between the groups ranged from 0.03% to 24.5%. Many variables had standardized differences <10% even before applying the IPTWs. After applying the IPTWs, all covariates had standardized differences <1.2%. Examination of kernel density plots confirmed that propensity scores for the groups overlapped, and no trimming was necessary.

Table 3 shows the crude rates and adjusted relative risks (aRRs) for each of the birth outcomes in the exposed and unexposed groups. Receiving HBPB was associated with reductions in low birth weight, preterm and small for gestational age births, and increases in breastfeeding initiation and large for gestational age births. HBPB was not associated with 5-minute Apgar scores or neonatal hospital readmissions. Table 3 also shows the mean birth hospitalization LOS for infants born to mothers exposed or unexposed to HBPB. For infants born by cesarean

delivery, there were no significant differences between groups ( $P = .87$ ); for infants born vaginally, receipt of HBPB was associated with shorter LOS ( $P < .0001$ ).

Sensitivity analyses found that HBPB's associations with breastfeeding initiation, low birth weight, preterm birth, and mean LOS for vaginal births were robust to unmeasured confounding (Table 3). After adjusting for the confounders included in the propensity score, there would need to be an unmeasured confounder that both perfectly predicted receipt of HBPB and accounted for ~60% of the relationship between HBPB and these 4 outcomes. The likelihood of such a confounder existing, after adjusting for covariates included in the propensity score, is very small. The findings regarding large for gestational age may be more sensitive to unmeasured confounding. Finally, the findings related to small for gestational age were very sensitive to unmeasured confounding and could potentially become nonsignificant if unmeasured confounders were included in our models.

Figure 2 illustrates the PAF for breastfeeding (4% increase) and PPF for low birth weight and preterm birth (21% and 17.5% decrease, respectively).



**FIGURE 2**  
PAF and PPF associated with receipt of the HBPB. GA, gestational age.

## DISCUSSION

Receipt of an unconditional prenatal income supplement by very-low-income women was associated with a number of positive outcomes: increased breastfeeding initiation; reductions in low birth weight, preterm births, and small for gestational age births; and shorter birth hospital stays for infants born vaginally. The provision of money to recipients without applying conditions differentiates this study from others in the literature. Birth outcomes improved, without requiring any specific actions from recipients to receive the income benefit or providing them with vouchers to buy specific food items.

As a society, we tend to assume that poor people cannot be trusted to make good choices. Indeed, when HBPB was first introduced in Manitoba, concerns were expressed about introducing a program for low-income women without conditions or accountability.<sup>55</sup> Although information about prenatal and infant health is included with the monthly payment, the Manitoba HBPB program trusts low-income women to make good choices regarding their pregnancies. There

is a growing body of evidence demonstrating improvements to child outcomes associated with increased family income that warrant attention from decision-makers.<sup>56–63</sup>

One strength of this study was the use of administrative data to identify all those eligible for HBPB, including those receiving and not receiving the supplement. Combining population-based databases on program participation with information on family risk factors and service use avoids the problems associated with reporting and recall bias.<sup>27</sup> Previous evaluations of similar programs have been limited by potential underreporting of program involvement and have proposed using administrative data to overcome this bias.

A further strength of this study was the availability of an extensive array of risk factors that are rarely available in administrative data.<sup>27,64</sup> These factors allowed us to balance measured differences between HBPB recipients and nonrecipients and ensure that those among our population of very-low-income women who received HBPB were comparable to those who did not, based on these observed

characteristics. Although we could not directly test whether the propensity score controlled for all unmeasured confounding, our analysis allowed us to assess the sensitivity of our findings.<sup>53</sup> We found that 4 of our 6 statistically significant associations were robust to unmeasured confounding: increased breastfeeding initiation and decreased low birth weight births, preterm births, and birth hospital LOS. Thus, based on the measured associations and the sensitivity analyses, the HBPB program seems to be improving these 4 infant health outcomes.

The benefits of breastfeeding, for both the developing infant<sup>65–69</sup> and the mother,<sup>70–74</sup> have been so clearly demonstrated that the US Surgeon General has called for action to support breastfeeding.<sup>75</sup> In our study, the increased breastfeeding initiation associated with HBPB was likely the result of information about the importance of breastfeeding sent with the monthly payment. The increase in breastfeeding is an important finding given that this very-low-income population is the least likely to breastfeed<sup>76</sup>; however, the PAF for breastfeeding was relatively small (4%), and our measure included only initiation, not duration. Research on similar programs (eg, WIC) has yielded equivocal results, with some studies actually finding reductions in breastfeeding initiation and duration associated with the program<sup>26,77–79</sup>; others that have more adequately controlled for confounding have found no association between WIC and breastfeeding.<sup>80</sup>

There is extensive literature on both the short- and long-term adverse effects of low birth weight<sup>2,5,7, 10,81–84</sup> and preterm births,<sup>13,85–87</sup> underscoring the importance of reducing these outcomes, particularly for vulnerable populations. Although reductions in low birth weight and preterm births have been found for

WIC in previous evaluations,<sup>24,29</sup> more sophisticated analyses suggest that previous positive findings may have more to do with selection bias than actual program effects.<sup>27,88</sup> Increases in birth weight were associated with WIC according to a design that exploited county-level variation in roll-out.<sup>29</sup> We found that the reductions in low birth weight and preterm births associated with HBPB were robust to unmeasured confounding, and they translated into the prevention of 21% of all low birth weight births and 17.5% of all preterm births for this vulnerable population.

Shorter hospital stays for uncomplicated vaginal births have medical, economic, and social benefits,<sup>89,90</sup> and they are indicative of better overall health of the mother and newborn.<sup>91</sup> We found that infants born to mothers receiving HBPB had significantly shorter birth hospital stays. Although modest, the LOS reduction associated with HBPB could translate into considerable savings in hospital days if this entire population received HBPB.

Despite the extensive array of risk factors available for propensity scores, a limitation of this study is the endogeneity bias that cannot be accounted for. The sensitivity analysis conducted attempted to address this factor; however, we cannot know for certain how unmeasured confounding influenced our findings.

Furthermore, to ensure comparability of income between our exposed and unexposed groups, we limited our evaluation to women receiving welfare rather than examining all low-income women receiving the income supplement during pregnancy. This approach limits the generalizability of our findings, although the population we examined is more comparable to the very-low-income women

participating in the US WIC program, and thus the findings may be applicable to that population.

A further limitation was our inability to determine why HBPB made a difference: was the additional money used for more nutritious food? Was stress reduced because rent could be paid on time? Research on the Earned Income Tax Credit in the United States suggests that increased income to low wage earners results in better nutritional intake for women in general,<sup>63</sup> decreased smoking for pregnant women,<sup>60,62,92</sup> and more prenatal care. Findings of increased infant birth weight associated with the Food Stamp Program also suggest that prenatal nutritional intake plays a role.<sup>93</sup> Future research should include qualitative analyses to explore how and why the modest monthly income supplement provided through HBPB made a difference to recipients' pregnancies. It is also important to explore the barriers that prevent eligible women from receiving HBPB.

## CONCLUSIONS

Using a quasi-experimental, retrospective cohort study design, we found that receipt of an unconditional income supplement by very-low-income women during pregnancy was associated with several positive outcomes: increased breastfeeding initiation, reductions in low birth weight and preterm births, and shorter mean length of birth hospital stay. Placing conditions on income supplements may not be necessary to promote prenatal and perinatal health.

## ACKNOWLEDGMENTS

This study was part of a program of research being conducted by the PATHS Equity Team: James Bolton, Marni Brownell, Charles Burchill, Elaine Burland, Mariette Chartier,

Dan Chateau, Malcolm Doupe, Greg Finlayson, Randall Fransoo, Chun Yan Goh, Milton Hu, Doug Jutte, Alan Katz, Laurence Katz, Lisa Lix, Patricia J. Martens (deceased), Colleen Metge, Nathan C. Nickel, Colette Raymond, Les Roos, Noralou Roos, Rob Santos, Joykrishna Sarkar, Mark Smith, Carole Taylor, and Randy Walld.

The authors acknowledge Jon Fischer for conducting a background literature search for the manuscript, Chun Yan Goh for preparation of tables and figures, and the following for providing input about the Healthy Baby Program: Shannon Dennehy, Joanne Waskin, Jan Sanderson, Leanne Boyd, Tamara Hes, and Susan Tessler. The authors also acknowledge the Manitoba Centre for Health Policy for use of data contained in the Population Health Research Data Repository under project 2012-006 (HIPC #2011/2012-24B). Data used in this study are from the Population Health Research Data Repository housed at MCHP, University of Manitoba, and were derived from data provided by the following: Healthy Child Manitoba; Manitoba Education and Advanced Learning; Manitoba Health, Healthy Living and Seniors; Manitoba Jobs and the Economy; and Statistics Canada.

## ABBREVIATIONS

aRR: adjusted relative risk  
CI: confidence interval  
HBPB: Healthy Baby Prenatal Benefit  
IPTW: inverse probability of treatment weights  
LOS: length of stay  
PAF: population-attributable fraction  
PPF: population-preventable fraction  
WIC: Special Supplemental Nutrition Program for Women, Infants, and Children

Mr Sarkar and Ms Taylor conducted the analyses, and reviewed and revised the manuscript; and Drs Brownell, Chartier, Nickel, Chateau, Burland, Jutte, Santos, Katz, Mr Sarkar, and Ms Taylor approved the final manuscript as submitted.

The results and conclusions are those of the authors, and no official endorsement by the Manitoba Centre for Health Policy, Manitoba Health, Healthy Living and Seniors, or other data providers is intended or should be inferred.

**DOI:** 10.1542/peds.2015-2992

Accepted for publication Mar 1, 2016

Address correspondence to Marni D. Brownell, PhD, 408-727 McDermot Ave, Winnipeg, MB, R3E 3P5, Canada. E-mail: marni\_brownell@cpe.umanitoba.ca

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** Dr Brownell acknowledges the financial support of the Government of Manitoba through the Manitoba Centre for Health Policy Population-Based Child Health Research Award. Dr Martens was supported by the Canadian Institutes of Health Research and the Public Health Agency of Canada with a CIHR/PHAC Applied Public Health Research Chair (2008–2014) during this research. Dr Katz acknowledges the support of the Manitoba Health Research Council and the Heart and Stroke Foundation for his Research Chair in Primary Prevention (2013–2018). None of the funders was involved in the: design of the study; conduct of the study; collection, management, analysis, or interpretation of the data; or the preparation, review, or approval of the manuscript. The other authors have indicated they have no financial relationships relevant to this article to disclose.

**FUNDING:** Supported by the Canadian Institutes of Health Research (grant number ROH-115206) and the Heart & Stroke Foundation of Canada (grant number PG-12-0534), under the program of research entitled “PATHS Equity for Children: A Program of Research into What Works to Reduce the Gap for Manitoba’s Children.”

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

## REFERENCES

1. Barker DJ. The origins of the developmental origins theory. *J Intern Med*. 2007;261(5):412–417
2. Jefferis BJ, Power C, Hertzman C. Birth weight, childhood socioeconomic environment, and cognitive development in the 1958 British birth cohort study. *BMJ*. 2002;325(7359):305
3. Wadhwa PD, Sandman CA, Porto M, Dunkel-Schetter C, Garite TJ. The association between prenatal stress and infant birth weight and gestational age at birth: a prospective investigation. *Am J Obstet Gynecol*. 1993;169(4):858–865
4. Finch BK. Socioeconomic gradients and low birth-weight: empirical and policy considerations. *Health Serv Res*. 2003;38(6 Pt 2):1819–1841
5. Kramer MS. Determinants of low birth weight: methodological assessment and meta-analysis. *Bull World Health Organ*. 1987;65(5):663–737
6. Kramer MS, Olivier M, McLean FH, Willis DM, Usher RH. Impact of intrauterine growth retardation and body proportionality on fetal and neonatal outcome. *Pediatrics*. 1990;86(5):707–713
7. Hack M, Klein NK, Taylor HG. Long-term developmental outcomes of low birth weight infants. *Future Child*. 1995;5(1):176–196
8. Mick E, Biederman J, Faraone SV, Sayer J, Kleinman S. Case-control study of attention-deficit hyperactivity disorder and maternal smoking, alcohol use, and drug use during pregnancy. *J Am Acad Child Adolesc Psychiatry*. 2002;41(4):378–385
9. Breslau N, Johnson EO, Lucia VC. Academic achievement of low birthweight children at age 11: the role of cognitive abilities at school entry. *J Abnorm Child Psychol*. 2001;29(4):273–279
10. Stein RE, Siegel MJ, Bauman LJ. Are children of moderately low birth weight at increased risk for poor health? A new look at an old question. *Pediatrics*. 2006;118(1):217–223
11. Ramsay MC, Reynolds CR. Does smoking by pregnant women influence IQ, birth weight, and developmental disabilities in their infants? A methodological review and multivariate analysis. *Neuropsychol Rev*. 2000;10(1):1–40
12. Lawlor DA, Batty GD, Morton SM, et al. Early life predictors of childhood intelligence: evidence from the Aberdeen children of the 1950s study. *J Epidemiol Community Health*. 2005;59(8):656–663
13. Huddy CL, Johnson A, Hope PL. Educational and behavioural problems in babies of 32-35 weeks gestation. *Arch Dis Child Fetal Neonatal Ed*. 2001;85(1):F23–F28
14. Rasmussen C, Horne K, Witol A. Neurobehavioral functioning in children with fetal alcohol spectrum disorder. *Child Neuropsychol*. 2006;12(6):453–468
15. Nigg JT, Breslau N. Prenatal smoking exposure, low birth weight, and disruptive behavior disorders. *J Am Acad Child Adolesc Psychiatry*. 2007;46(3):362–369
16. Lu MC, Chen B. Racial and ethnic disparities in preterm birth: the role of stressful life events. *Am J Obstet Gynecol*. 2004;191(3):691–699
17. Lu MC, Kotelchuck M, Hogan V, Jones L, Wright K, Halfon N. Closing the black-white gap in birth outcomes: a life-course approach. *Ethn Dis*. 2010;20(1 suppl 2):S2–S62, 76
18. Blumenshine P, Egarter S, Barclay CJ, Cubbin C, Braveman PA. Socioeconomic disparities in adverse birth outcomes: a systematic review. *Am J Prev Med*. 2010;39(3):263–272
19. Fernald LC, Gertler PJ, Neufeld LM. Role of cash in conditional cash transfer programmes for child health, growth, and development: an analysis of Mexico's Oportunidades. *Lancet*. 2008;371(9615):828–837
20. Paes-Sousa R, Santos LM, Miazaki ES. Effects of a conditional cash transfer

- programme on child nutrition in Brazil. *Bull World Health Organ*. 2011;89(7):496–503
21. Lagarde M, Haines A, Palmer N. The impact of conditional cash transfers on health outcomes and use of health services in low and middle income countries. *Cochrane Database Syst Rev*. 2009;(4):CD008137
22. Barber SL, Gertler PJ. The impact of Mexico's conditional cash transfer programme, Oportunidades, on birthweight. *Trop Med Int Health*. 2008;13(11):1405–1414
23. USDA Food and Nutrition Service. Women, Infants and Children (WIC). Available at: <http://www.fns.usda.gov/wic/about-wic-wic-glance>. Accessed February 28, 2014
24. Abrams B. Preventing low birth weight: does WIC work? A review of evaluations of the Special Supplemental Food Program for Women, Infants, and Children. *Ann N Y Acad Sci*. 1993;678:306–316
25. Avruch S, Cackley AP. Savings achieved by giving WIC benefits to women prenatally. *Public Health Rep*. 1995;110(1):27–34
26. Bitler MP, Currie J. Does WIC work? The effects of WIC on pregnancy and birth outcomes. *J Policy Anal Manage*. 2005;24(1):73–91
27. Foster EM, Jiang M, Gibson-Davis CM. The effect of the WIC program on the health of newborns. *Health Serv Res*. 2010;45(4):1083–1104
28. Kowaleski-Jones L, Duncan GJ. Effects of participation in the WIC program on birthweight: evidence from the National Longitudinal Survey of Youth. Special Supplemental Nutrition Program for Women, Infants, and Children. *Am J Public Health*. 2002;92(5):799–804
29. Hoynes H, Page M, Stevens AH. Can targeted transfers improve birth outcomes? Evidence from the introduction of the WIC program. *J Public Econ*. 2011;95(7-8):813–827
30. Nickel NC, Chateau DG, Martens PJ, et al; PATHS Equity Team. Data resource profile: Pathways to Health and Social Equity for Children (PATHS Equity for Children). *Int J Epidemiol*. 2014;43(5):1438–1449
31. Roos NP, Roos LL, Brownell M, Fuller EL. Enhancing policymakers' understanding of disparities: relevant data from an information-rich environment. *Milbank Q*. 2010;88(3):382–403
32. Roos LL, Nicol JP. A research registry: uses, development, and accuracy. *J Clin Epidemiol*. 1999;52(1):39–47
33. Roos LL, Brownell M, Lix L, Roos NP, Walld R, MacWilliam L. From health research to social research: privacy, methods, approaches. *Soc Sci Med*. 2008;66(1):117–129
34. Brownell MD, Chartier M, Santos R, Au W, Roos NP, Girard D. Evaluation of a newborn screen for predicting out-of-home placement. *Child Maltreat*. 2011;16(4):239–249
35. Roos NP, Brownell M, Guevremont A, et al. The complete story: a population-based perspective on school performance and educational testing. *Can J Educ*. 2006;29(3):684–705
36. Roos LL Jr, Nicol JP, Cageorge SM. Using administrative data for longitudinal research: comparisons with primary data collection. *J Chronic Dis*. 1987;40(1):41–49
37. Roos LL, Menec V, Currie RJ. Policy analysis in an information-rich environment. *Soc Sci Med*. 2004;58(11):2231–2241
38. Roos LL, Gupta S, Soodeen RA, Jebamani L. Data quality in an information-rich environment: Canada as an example. *Can J Aging*. 2005;24(suppl 1):153–170
39. Robinson JR, Young TK, Roos LL, Gelskey DE. Estimating the burden of disease. Comparing administrative data and self-reports. *Med Care*. 1997;35(9):932–947
40. Oreopoulos P, Stabile M, Walld R, Roos LL. Short-, medium-, and long-term consequences of poor infant health: an analysis using siblings and twins. *J Hum Resour*. 2008;43:88–138
41. Heaman M, Kingston D, Helewa M, et al. Perinatal services and outcomes in Manitoba. Available at: [http://mchp-appserv.cpe.umanitoba.ca/reference/perinatal\\_report\\_WEB.pdf](http://mchp-appserv.cpe.umanitoba.ca/reference/perinatal_report_WEB.pdf). Accessed July 29, 2015
42. Brownell M, Chartier M, Au W, Schultz J. Evaluation of the Healthy Baby Program. Available at: [http://mchp-appserv.cpe.umanitoba.ca/reference/MCHP-Healthy\\_Baby\\_Full\\_Report\\_WEB.pdf](http://mchp-appserv.cpe.umanitoba.ca/reference/MCHP-Healthy_Baby_Full_Report_WEB.pdf). Accessed March 11, 2015
43. Kramer MS, Platt RW, Wen SW, et al; Fetal/Infant Health Study Group of the Canadian Perinatal Surveillance System. A new and improved population-based Canadian reference for birth weight for gestational age. *Pediatrics*. 2001;108(2):E35
44. Lix L, Yogendran M, Burchill C, et al. Defining and validating chronic diseases: an administrative data approach. Winnipeg, Canada: Manitoba Centre for Health Policy; 2006
45. Ruth CA, Roos NP, Hildes-Ripstein E, Brownell MD. Infants born to mothers with diabetes in pregnancy at the population level in Manitoba: more questions than answers. *Can J Diabetes*. 2012;36(2):71–74
46. Jutte DP, Roos NP, Brownell MD, Briggs G, MacWilliam L, Roos LL. The ripples of adolescent motherhood: social, educational, and medical outcomes for children of teen and prior teen mothers. *Acad Pediatr*. 2010;10(5):293–301
47. Chateau D, Metge C, Prior H, Soodeen RA. Learning from the census: the socio-economic factor index (SEFI) and health outcomes in Manitoba. *Can J Public Health*. 2012;103(suppl 8):S23–S27
48. Foster EM. Causal inference and developmental psychology. *Dev Psychol*. 2010;46(6):1454–1480
49. Rubin D. Using propensity scores to help design observational studies: application to the tobacco litigation. *Health Serv Outcomes Res Methodol*. 2001;2:169–188
50. Austin PC. Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples. *Stat Med*. 2009;28(25):3083–3107
51. Guo SY, Fraser MW. *Propensity Score Analysis: Statistical Methods and Applications*. Thousand Oaks, CA: SAGE Publications, Inc.; 2010
52. Morgan SL, Winship C. *Counterfactuals and Causal Inference: Methods and Principles for Social Research*. New

- York, NY: Cambridge University Press; 2007
53. Rosenbaum PR. *Observational Studies (Springer Series in Statistics)*, 2nd ed. New York, NY: Springer-Verlag New York, Inc; 2010
54. Rockhill B, Newman B, Weinberg C. Use and misuse of population attributable fractions. *Am J Public Health*. 1998;88(1):15–19
55. Legislative Assembly of Manitoba. Available at: [http://gov.mb.ca/legislature/hansard/37th\\_2nd/vol\\_014/h014.html](http://gov.mb.ca/legislature/hansard/37th_2nd/vol_014/h014.html). Accessed December 21, 2014
56. Akee RK, Copeland WE, Keeler G, Angold A, Costello EJ. Parents' incomes and children's outcomes: a quasi-experiment. *Am Econ J Appl Econ*. 2010;2(1):86–115
57. Costello EJ, Compton SN, Keeler G, Angold A. Relationships between poverty and psychopathology: a natural experiment. *JAMA*. 2003;290(15):2023–2029
58. Costello EJ, Erkanli A, Copeland W, Angold A. Association of family income supplements in adolescence with development of psychiatric and substance use disorders in adulthood among an American Indian population. *JAMA*. 2010;303(19):1954–1960
59. Forget EL. The town with no poverty: the health effects of a Canadian Guaranteed Annual Income field experiment. *Can Public Policy*. 2011;37(3):283–305
60. Hamad R, Rehkopf DH. Poverty, pregnancy, and birth outcomes: a study of the Earned Income Tax Credit. *Paediatr Perinat Epidemiol*. 2015;29(5):444–452
61. Paxson C, Schady N. Does money matter? The effects of cash transfers on child development in rural Ecuador. *Econ Dev Cult Change*. 2010;59(1):187–229
62. Rehkopf DH, Strully KW, Dow WH. The short-term impacts of Earned Income Tax Credit disbursement on health. *Int J Epidemiol*. 2014;43(6):1884–1894
63. Strully KW, Rehkopf DH, Xuan Z. Effects of prenatal poverty on infant health: state earned income tax credits and birth weight. *Am Sociol Rev*. 2010;75(4):534–562
64. Meghea CI, Raffo JE, VanderMeulen P, Roman LA. Moving toward evidence-based federal Healthy Start program evaluations: accounting for bias in birth outcomes studies. *Am J Public Health*. 2014;104(suppl 1):S25–S27
65. Quigley MA, Hockley C, Carson C, Kelly Y, Renfrew MJ, Sacker A. Breastfeeding is associated with improved child cognitive development: a population-based cohort study. *J Pediatr*. 2012;160(1):25–32
66. McCrory C, Layte R. Breastfeeding and risk of overweight and obesity at nine-years of age. *Soc Sci Med*. 2012;75(2):323–330
67. Kramer MS, Kakuma R. Optimal duration of exclusive breastfeeding. *Cochrane Database Syst Rev*. 2002;(1):CD003517
68. Ip S, Chung M, Raman G, et al. Breastfeeding and maternal and infant health outcomes in developed countries. *Evid Rep Technol Assess (Full Rep)*. 2007;(153):1–186
69. Heikkilä K, Sacker A, Kelly Y, Renfrew MJ, Quigley MA. Breast feeding and child behaviour in the Millennium Cohort Study. *Arch Dis Child*. 2011;96(7):635–642
70. Cramer DW. The epidemiology of endometrial and ovarian cancer. *Hematol Oncol Clin North Am*. 2012;26(1):1–12
71. Kobayashi S, Sugiyama H, Ando Y, et al. Reproductive history and breast cancer risk. *Breast Cancer*. 2012;19(4):302–308
72. Taylor JS, Kacmar JE, Nothnagle M, Lawrence RA. A systematic review of the literature associating breastfeeding with type 2 diabetes and gestational diabetes. *J Am Coll Nutr*. 2005;24(5):320–326
73. Tharner A, Luijk MP, Raat H, et al. Breastfeeding and its relation to maternal sensitivity and infant attachment. *J Dev Behav Pediatr*. 2012;33(5):396–404
74. Ystrom E. Breastfeeding cessation and symptoms of anxiety and depression: a longitudinal cohort study. *BMC Pregnancy Childbirth*. 2012;12:36
75. US Department of Health and Human Services, Office of the Surgeon General. The Surgeon General's Call to Action to Support Breastfeeding 2011. Available at: <http://www.surgeongeneral.gov/library/calls/breastfeeding/calltoactiontosupportbreastfeeding.pdf>. Accessed February 1, 2012
76. Li R, Darling N, Maurice E, Barker L, Grummer-Strawn LM. Breastfeeding rates in the United States by characteristics of the child, mother, or family: the 2002 National Immunization Survey. *Pediatrics*. 2005;115(1). Available at: [www.pediatrics.org/cgi/content/full/115/1/e31](http://www.pediatrics.org/cgi/content/full/115/1/e31)
77. Chatterji P, Brooks-Gunn J. WIC participation, breastfeeding practices, and well-child care among unmarried, low-income mothers. *Am J Public Health*. 2004;94(8):1324–1327
78. Jacknowitz A, Novillo D, Tihen L. Special Supplemental Nutrition Program for Women, Infants, and Children and infant feeding practices. *Pediatrics*. 2007;119(2):281–289
79. Ziol-Guest KM, Hernandez DC. First- and second-trimester WIC participation is associated with lower rates of breastfeeding and early introduction of cow's milk during infancy. *J Am Diet Assoc*. 2010;110(5):702–709
80. Jiang M, Foster EM, Gibson-Davis CM. The effect of WIC on breastfeeding: a new look at an established relationship. *Child Youth Serv Rev*. 2010;32(2):264–273
81. McCormack VA, dos Santos Silva I, Koupil I, Leon DA, Lithell HO. Birth characteristics and adult cancer incidence: Swedish cohort of over 11, 000 men and women. *Int J Cancer*. 2005;115(4):611–617
82. Frankel S, Elwood P, Sweetnam P, Yarnell J, Smith GD. Birthweight, body-mass index in middle age, and incident coronary heart disease. *Lancet*. 1996;348(9040):1478–1480
83. Leon DA, Lithell HO, Vågerö D, et al. Reduced fetal growth rate and increased risk of death from ischaemic heart disease: cohort study of 15 000 Swedish men and women born 1915–29. *BMJ*. 1998;317(7153):241–245
84. Forsén T, Eriksson J, Tuomilehto J, Reunanen A, Osmond C, Barker D. The fetal and childhood growth of persons who develop type 2 diabetes. *Ann Intern Med*. 2000;133(3):176–182

85. Goldenberg RL. The management of preterm labor. *Obstet Gynecol.* 2002;100(5 Pt 1):1020–1037
86. Health Canada. Canadian Perinatal Health Report, 2003. Available at: <http://publications.gc.ca/collections/Collection/H49-142-2003E.pdf>. Accessed March 17, 2010
87. Mathews TJ, Menacker F, MacDorman MF. Infant mortality statistics from the 2001 period linked birth/infant death data set. *Natl Vital Stat Rep.* 2003;52(2):1–28
88. Joyce T, Racine A, Yunzal-Butler C. Reassessing the WIC effect: evidence from the Pregnancy Nutrition Surveillance System. *J Policy Anal Manage.* 2008;27(2):277–303
89. Danielsen B, Castles AG, Damberg CL, Gould JB. Newborn discharge timing and readmissions: California, 1992-1995. *Pediatrics.* 2000;106(1 pt 1):31–39
90. Lee KS, Perlman M, Ballantyne M, Elliott I, To T. Association between duration of neonatal hospital stay and readmission rate. *J Pediatr.* 1995;127(5):758–766
91. American Academy of Pediatrics. Committee on Fetus and Newborn. Hospital stay for healthy term newborns. *Pediatrics.* 2010;125(2):405–409
92. Hoynes H, Miller D, Simon D. Income, the Earned Income Tax Credit, and infant health. *Am Econ J Econ Policy.* 2015;7(1):172–211
93. Almond D, Hoynes HW, Schanzenbach DW. Inside the war on poverty: the impact of food stamps on birth outcomes. *Rev Econ Stat.* 2011;93(2):387–403

## Unconditional Prenatal Income Supplement and Birth Outcomes

Marni D. Brownell, Mariette J. Chartier, Nathan C. Nickel, Dan Chateau, Patricia J. Martens, Joykrishna Sarkar, Elaine Burland, Douglas P. Jutte, Carole Taylor, Robert G. Santos, Alan Katz and On behalf of the PATHS Equity for Children Team

*Pediatrics* 2016;137;; originally published online May 12, 2016;

DOI: 10.1542/peds.2015-2992

### Updated Information & Services

including high resolution figures, can be found at:  
</content/137/6/e20152992.full.html>

### References

This article cites 81 articles, 15 of which can be accessed free at:  
</content/137/6/e20152992.full.html#ref-list-1>

### Subspecialty Collections

This article, along with others on similar topics, appears in the following collection(s):

**Fetus/Newborn Infant**

[/cgi/collection/fetus:newborn\\_infant\\_sub](/cgi/collection/fetus:newborn_infant_sub)

**Public Health**

[/cgi/collection/public\\_health\\_sub](/cgi/collection/public_health_sub)

### Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:  
</site/misc/Permissions.xhtml>

### Reprints

Information about ordering reprints can be found online:  
</site/misc/reprints.xhtml>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Unconditional Prenatal Income Supplement and Birth Outcomes**

Marni D. Brownell, Mariette J. Chartier, Nathan C. Nickel, Dan Chateau, Patricia J. Martens, Joykrishna Sarkar, Elaine Burland, Douglas P. Jutte, Carole Taylor, Robert G. Santos, Alan Katz and On behalf of the PATHS Equity for Children Team

*Pediatrics* 2016;137;; originally published online May 12, 2016;

DOI: 10.1542/peds.2015-2992

The online version of this article, along with updated information and services, is located on the World Wide Web at:

</content/137/6/e20152992.full.html>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2016 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

