


Association Studies



Overview

- Association Studies
 - Design
 - Analysis
- Candidate Gene Studies
 - Resources
 - Selecting 'tag' SNPs
- Pathways


Process of Genetic Epidemiology



Association Studies

Approaches to Identifying Susceptibility Genes for Common Diseases


Linkage Studies:




- A relation between loci
- Done within families

Association Studies:

- A relation between alleles
- Done in populations



Association Studies



Association Studies

- Use of (population-based) association studies is rapidly expanding, reflecting a number of good properties, including:
 - Ease, since one need not collect large pedigrees
 - Potential for being more powerful than conventional linkage-based approaches
- Common disease / common variant hypothesis

Linkage vs. Association

Linkage					Association			
Genotypic risk ratio (γ)	Frequency of disease allele A (p)	Probability of allele sharing (Y)	No. of families required (N)	Probability of transmitting disease allele A $P(\text{tr-A})$	Singletons		Sib pairs	
					Proportion of heterozygous parents (Het)	(N)	(Het)	(N)
4.0	0.01	0.520	4260	0.800	0.048	1098	0.112	235
	0.10	0.597	185	0.800	0.346	150	0.537	48
	0.50	0.576	297	0.800	0.500	103	0.424	61
	0.80	0.529	2013	0.800	0.235	222	0.163	161
2.0	0.01	0.502	296,710	0.667	0.029	5823	0.043	1970
	0.10	0.518	5382	0.667	0.245	695	0.323	264
	0.50	0.526	2498	0.667	0.500	340	0.474	180
	0.80	0.512	11,917	0.667	0.267	640	0.217	394
1.5	0.01	0.501	4,620,807	0.600	0.025	19,320	0.031	7776
	0.10	0.505	67,816	0.600	0.197	2218	0.253	941
	0.50	0.510	17,997	0.600	0.500	949	0.490	484
	0.80	0.505	67,816	0.600	0.286	1663	0.253	941

Comparison of linkage and association studies. Number of families needed for identification of a disease gene.

Risch & Merikangas, Science 1996


Association Studies – Educational Group Exercise!

- Come up with important aspects and determine the design AND how to analyse your genotype association study...
 - What aspects might influence where you draw your cases/controls?
 - How would you choose where in the genome?
 - Do you need all of that? Is there a way you could genotype less?
 - How might you conduct a test of if the allele frequencies differ in cases and controls at a specific SNP?




Not about getting things perfectly right, the exercise is to get you to think about the problem, as studies show that you will remember things better that way.

Association Studies



Hirschhorn & Daly, Nat Rev Genet 2005



Key concept: Linkage Disequilibrium

Implications on design?
Implications on analysis?

Association Study Approaches

- Direct vs Indirect (tagSNPs, later in lecture)
- Candidate genes:
 - Functional
 - All common variants
 - Exome Arrays
- All common variants in genome (GWAS)
- All variants in genes/genome (sequencing)
 - Expensive

Genomics Revolution

Human Genome Project:

13 years, \$3B for 1 sequence

Now: <1 week, ~\$1.5K


> 500 times faster

< one millionth the cost!

Still not quite there:

1 hour, \$1K

(#1 Innovation, 2010)




The Economist, 2010

Study Design: Control Selection

- A critical aspect of association studies is that controls should be selected from the cases' source population.
- That is, controls should be those individuals who, if they were diseased, would become cases.

Family-Based Association Studies

(Covered in last lecture)



Termed “**pseudo**”-controls (the hypothetical other mendelian transmissions)


Continuum of Assoc Study Designs



Population Stratification

Overmatching

(Bias (**maybe**).....versus..... less efficient)



Association Analysis

Genotype	Cases	Controls	OR
TT	A	D	1
CT	B	E	BD/AE
CC	C	F	CD/AF

Simple chi-square test comparing genotype frequencies (2 d.f.)
Called a co-dominant analysis

Simple chi-squared test

Observed:

Geno	Case	Control	Total
TT	A	D	A+D=nCC
CT	B	E	B+E=nCT
CC	C	F	C+F=nTT
Total	A+B+C	D+E+F	A+B+C+D+E+F
	=nCase	=nCont	=n

Expected

Case	Control
$n_{TT} * n_{Case} / n$	$n_{TT} * n_{Cont} / n$
$n_{CT} * n_{Case} / n$	$n_{CT} * n_{Cont} / n$
$n_{CC} * n_{Case} / n$	$n_{CC} * n_{Cont} / n$

Sum $(\text{Observed} - \text{Expected})^2 / \text{Expected}$ as before. Chi squared with 2 degrees of freedom.

Hint: Expected cell count = row_total * column_total / total

Simple chi-squared test example

Observed:				Expected		
Geno	Case	Control	Total	Case	Control	OR (to TT genotype)
TT	5	15	20	$20 \times 35 / 65 = 10.8$	$20 \times 30 / 65 = 9.2$	1
CT	10	10	20	$20 \times 35 / 65 = 10.8$	$20 \times 30 / 65 = 9.2$	$(10 \times 15) / (5 \times 10) = 3$
CC	20	5	25	$25 \times 35 / 65 = 13.5$	$25 \times 30 / 65 = 11.5$	$(20 \times 15) / (5 \times 5) = 12$
Total	35	30	65			
	=nCase	=nCont	=n			

Sum $(\text{Observed} - \text{Expected})^2 / \text{Expected}$

$$\begin{aligned}
 &= (5 - 10.8)^2 / 10.8 + (10 - 10.8)^2 / 10.8 + (20 - 13.5)^2 / 13.5 \\
 &\quad + (15 - 9.2)^2 / 9.2 + (10 - 9.2)^2 / 9.2 + (5 - 11.5)^2 / 11.5 \\
 &= 13.7
 \end{aligned}$$

P-value = 0.0011

Genetic Model

Genotype	OR	ORs depend on genetic model
TT	1	$R = r = 1$ not risk allele
CT	r	$R > r = 1$ recessive
CC	R	$R = r > 1$ dominant
		$R = r^2 > 1$ log additive

(Assuming positive association)

Tests of association

If genetic model known:

- Collapse genotypes into 2x2 table, 1 d.f. test
- Trend test for log additive
- Use logistic regression: coding; covariates

- Rarely know genetic model
- Use all three models (dom, rec, log additive)
- **Compare fit with the co-dominant (2d.f.) model (LR test)** / it's two estimated coefficients
- Cannot use LR test to compare models with each other as not nested
- Model with best fit and smallest P is best?

Tests of Association

2 df Genotype			Recessive (G)			Dominant (G)		
Genotype	Case	Control	Case Control			Case Control		
GG	A	D	GG	A	D	GG or GT	A+B	D+E
GT	B	E	GT or TT	B+C	E+F	TT	C	F
TT	C	F						
~chi_sq(2df)			~chi_sq(1df)			~chi_sq(1df)		
Genotype	Case	Control	Case Control			Case Control		
TT	5	15	TT	5	15	TT or TG	15	25
GG	20	5	GG or GT	30	15	GG	20	5
GT	10	10						
P=0.0011			P=0.0045			P=0.0020		

Tests of Association – OR use logistic regression, covariates...

```
. * Load in the dataset, and describe it
```

```
.  
. cd c:\ttemp  
c:\ttemp
```

```
.  
. insheet using geno_ex.csv, clear  
(2 vars, 65 obs)
```

```
.  
. table geno_logadditive case
```

```
-----  
geno_log |      case  
additive |      0      1  
-----+-----  
      0 |     15     5  
      1 |     10    10  
      2 |      5    20  
-----
```

Tests of Association – OR use logistic regression, covariates...

```
. * handle a model fitting both coefficients
. generate geno1 = (geno_logadditive==1)
. generate geno2 = (geno_logadditive==2)
. logit case geno1 geno2
```

```
Iteration 0:  log likelihood = -44.862069
Iteration 1:  log likelihood = -37.630625
Iteration 2:  log likelihood = -37.619709
Iteration 3:  log likelihood = -37.619707
```

Logistic regression	Number of obs	=	65
	LR chi2(2)	=	14.48
	Prob > chi2	=	0.0007
Log likelihood = -37.619707	Pseudo R2	=	0.1614

case	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
geno1	1.098612	.68313	1.61	0.108	-.2402981	2.437522
geno2	2.484906	.7187953	3.46	0.001	1.076094	3.893719
_cons	-1.098612	.5163978	-2.13	0.033	-2.110733	-.0864911

Tests of Association – OR use logistic regression, covariates...

```
. * handle a model fitting both coefficients  
. generate geno1 = (geno_logadditive==1)  
. generate geno2 = (geno_logadditive==2)  
. logit case geno1 geno2
```

```
Iteration 0:  log likelihood = -44.862069  
Iteration 1:  log likelihood = -37.630625  
Iteration 2:  log likelihood = -37.619709  
Iteration 3:  log likelihood = -37.619707
```

```
Logistic regression               Number of obs   =          65  
                                LR chi2(2)         =          14.48  
                                Prob > chi2         =          0.0007  
Log likelihood = -37.619707       Pseudo R2        =          0.1614
```

case	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
-----+-----						
geno1	1.098612	.68313	1.61	0.108	-.2402981	2.437522
geno2	2.484906	.7187953	3.46	0.001	1.076094	3.893719
_cons	-1.098612	.5163978	-2.13	0.033	-2.110733	-.0864911
-----+-----						

**OR=exp(1.0986)
=3**

**OR=exp(2.4849)
=12**

**What model
does this
resemble?**

Tests of Association – OR use logistic regression, covariates...

```
. * recessive test for T / dominant test for G
. generate geno0 = (geno_logadditive==0)
. logit case geno0
```

```
Iteration 0:  log likelihood = -44.862069
Iteration 1:  log likelihood = -39.903582
Iteration 2:  log likelihood = -39.889842
Iteration 3:  log likelihood = -39.88984
```

Logistic regression	Number of obs	=	65
	LR chi2(1)	=	9.94
	Prob > chi2	=	0.0016
Log likelihood = -39.88984	Pseudo R2	=	0.1108

case	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]
-----+-----					
geno0	-1.791759	.6055301	-2.96	0.003	-2.978576 - .6049422
_cons	.6931472	.3162278	2.19	0.028	.0733521 1.312942

Tests of Association – OR use logistic regression, covariates...

```
. * dominant test for G / recessive test for T  
. logit case geno2
```

```
Iteration 0:  log likelihood = -44.862069  
Iteration 1:  log likelihood = -39.003467  
Iteration 2:  log likelihood = -38.972594  
Iteration 3:  log likelihood = -38.97259
```

Logistic regression	Number of obs	=	65
	LR chi2(1)	=	11.78
	Prob > chi2	=	0.0006
Log likelihood = -38.97259	Pseudo R2	=	0.1313

case	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]
geno2	1.89712	.5972157	3.18	0.001	.7265984 3.067641
_cons	-.5108255	.3265986	-1.56	0.118	-1.150947 .129296

Tests of Association – OR use logistic regression, covariates...

. * log additive

```
. logit case geno
```

Logistic regression

Number of obs = 65

$$\text{LR } \chi^2(1) = 14.42$$

```
Prob > chi2      = 0.00014
```

case	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
geno	1.2451	.3592	3.47	0.0005	.5360	1.9440
_cons	-1.1577	.4664	-2.48	0.0131	-2.0461	-0.2359

Review: Population stratification


Population stratification is a form of confounding in genetic studies where a gene under study shows marked variation in allele frequency across subgroups of a population and these subgroups differ in their baseline risk of disease

(Pima and Papago example, significance indicates what population from, not association with trait.)

Correcting for population stratification

- “Homogeneous” group (and ignore)
- Genomic control (PMID: 11315092) [see next slide]
 - Non-central chi-square λ = median of all χ^2 tests in the sample
 - Adjust all test statistics for inflation due to empirical chi-square distribution ($\chi^2_{\text{new}} = \chi^2_{\text{old}}/\lambda$)
 - Critiques: An average across the genome, and may **over** or under correct for individual tests
 - For large samples, e.g. 100,000, polygenetic traits, should **NOT** equal 1!
- Ancestry Principal Components (more in GWAS lecture)
 - Eigenstrat (PMID: 16862161)
 - Adjust regression model for principal component values which serve as a proxy for ancestry
- Mixed model – more in GWAS lecture (PMID 20562875) [idea is to condition on the rest of the genome]


Quantile-Quantile (QQ) Plots



3. Candidate Gene Studies

- Selection of candidates
Linkage regions?
Biological support?

“I am interested in a candidate gene and have samples ready to study. What SNPs do I genotype?”



Candidate Gene: Where do I Start?

- Location:

What chromosome? What position on the chr?


- Exons/UTR:

How many exons? UTR regions?

- Size:

How large is the gene?

Use UCSC genome browser.



SNP Picking: Things to Consider

- Validation: What is the quality of the SNPs?
- Informativity: Are these SNPs informative in my population?
How common are they? Location?
- Potentially Functional: Do these SNPs have a potential biological impact? Missense variants?
- Previously Associated: Have previous studies found SNPs in the candidate gene associated with the outcome?

Learning activity! Try these websites with these terms

- Links:
 - UCSC Genome browser / 1000 Genomes Browser
 - dbSNP
 - Pubmed
- Example terms: mthfr (a gene), mthfr cancer
- Things to find out: Chromosome? # Exons? # Introns? Size of gene? Functional variants? Previously associated variants? SNP validation status?


UCSC Genome Browser

UCSC Genome Browser on Human Dec. 2013 (GRCh38/hg38) Assembly

move <<< << < > >> >>> zoom in 1.5x 3x 10x base zoom out 1.5x 3x 10x 100x

chr1:11,786,603-11,806,058 19,456 bp. mthfr

go



SNP Picking: Validation

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?CMD=DisplayFiltered&DB=snp

dbSNP Go

Single Nucleotide Polymorphism (SNP)

NCBI

ENTREZ **SNP**
Single Nucleotide Polymorphism

PubMed Nucleotide Protein Genome

NP for Go Clear Save Search

Limits Preview/Index History Clipboard Details

Display Graphic Summary Show 20 Sort by Send to

All: 228 Human: 109 Mouse: 119 NEW: 107 Other Organisms: 0 UPDATE: 7

Items 1 - 20 of 109

☐ 1: [rs28484963](#) [*Homo sapiens*]

ACACTTTGCAATCCTGCCCCGCCCC [G/T] GTGATTTCTCTTTGGGCGGCAGGCA

MapView GeneView SeqView No 3D No OMIM

☐ 2: [rs17421560](#) [*Homo sapiens*]


ATTTTCTATCATGAACAGCTGAGG [A/G] CTTTCTTTTTTAAATGTTTTTATT

MapView GeneView SeqView No 3D No OMIM

☐ 3: [rs17421511](#) [*Homo sapiens*]

SNP Picking: Validation

Single Nucleotide Polymorphism



Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books	SNP
Search for SNP on NCBI Reference Assembly							
▼ for Go							
Reference SNP(refSNP) Cluster Report: rs28484963							
refSNP ID: rs28484963				Allele			
Organism: human (<i>Homo sapiens</i>)				Variation Class: SNP: single nucleotide polymorphism			
Molecule Type: Genomic				Alleles: G/T			
Created/Updated in build: 125/125				Ancestral Allele: T			
Map to Genome Build: 36.1							

SNP Details are organized in the following sections:

[Submission](#) [Fasta](#) [Resource](#) [GeneView](#) [Map](#) [Diversity](#) [Validation](#)

Submitter records for this RefSNP Cluster

The submission [ss35207915](#) has the longest flanking sequence of all cluster members and w

NCBI Assay ID	Handle Submitter ID	Validation Status	Orientation /Strand	Alleles	5' Near Seq 30 bp
ss35207915	SSAHASNP TA-079.chr1_11782008		fwd/B	G/T	agaaaacactttgcattcctgccc

Validation Summary:

Validation status	Marker displays Mendelian segregation	PCR results confirmed in multiple reactions	Homozygotes detected in individual genotype data
	UNKNOWN	UNKNOWN	UNKNOWN

SNP Picking: Informative

Population Diversity

Population Diversity

ss#	Sample Ascertainment				Genotype Detail ^{NEW}				Alleles	
	Population	Individual Group	Chrom. Sample Cnt.	Source	A/A	A/G	G/G	HWP	A	G
rs24246985	AFD_EUR_PANEL	European	48	IG	0.375	0.625	0.273	0.188	0.812	
	AFD_AFR_PANEL	African American	46	IG		1.000			1.000	
	AFD_CHN_PANEL	Asian	48	IG	0.042	0.958	1.000	0.021	0.979	
	HapMap-CEU	European	120	IG	0.050	0.317	0.633	1.000	0.208	0.792
	HapMap-HCB	Asian	90	IG	0.200	0.800	0.479	0.100	0.900	
	HapMap-JPT	Asian	90	IG	0.156	0.844	0.584	0.078	0.922	
	HapMap-YRI	Sub-Saharan African	120	IG		1.000			1.000	
	CHMJ		74	IG			0.001	0.041	0.959	

(rs17421511)

SNP Picking: Potentially Functional

Region	Contig position	mRNA pos	dbSNP rs# cluster id	Hetero-zygosity	Validation	3D	OMIM	Function	dbSNP allele	Protein residue	Codon pos	Amino acid pos
exon_12	6388294	1832	rs2274976	0.129	  H			nonsynonymous	A	Gln [Q]	2	594
				0.129	  H			contig reference	G	Arg [R]	2	594
exon_11	6388686	1748	rs2274974	N.D.		H		nonsynonymous	A	Glu [E]	2	566
				N.D.		H		contig reference	G	Gly [G]	2	566
exon_8	6391824	1356	rs4846051	0.164	  H			synonymous	T	Phe [F]	3	435
				0.164	  H			contig reference	C	Phe [F]	3	435
	6391843	1337	rs1801131	0.351	  H			nonsynonymous	C	Ala [A]	2	429
				0.351	  H	Yes		contig reference	A	Glu [E]	2	429
exon_7	6392263	1107	rs2066462	0.176	  H			synonymous	T	Ser [S]	3	352
				0.176	  H			contig reference	C	Ser [S]	3	352
exon_6	6392608	996	rs6664734	N.D.		H	Yes	synonymous	A	Val [V]	3	315
				N.D.		H	Yes	contig reference	G	Val [V]	3	315
	2647	957	rs13306555	N.D.		H	Yes	synonymous	T	Ala [A]	3	302
				N.D.		H	Yes	contig reference	C	Ala [A]	3	302
exon_5	6393745	716	rs1801133	0.403	  H	Yes		nonsynonymous	T	Val [V]	2	222
				0.403	  H	Yes	Yes	contig reference	C	Ala [A]	2	222
	6393804	657	rs13306554	N.D.	  H	Yes		synonymous	T	Pro [P]	3	202
				N.D.	  H	Yes		contig reference	C	Pro [P]	3	202
exon_3	6398643	468	rs2066466	0.050	  H	Yes		synonymous	A	Thr [T]	3	139
				0.050	  H	Yes		contig reference	G	Thr [T]	3	139
	6398715	396	rs2066461	0.026		H	Yes	synonymous	A	Thr [T]	3	115

SNP Picking: Previously Associated

www.ncbi.nlm.nih.gov/pubmed?term=mthfr cancer

Comics R TclTk Jobs Slashdot fbook News Articles PLINK New Tab UCSF THProject Google Sites uw e-mail

NCBI Resources How To

PubMed.gov
US National Library of Medicine
National Institutes of Health

PubMed mthfr cancer

RSS Save search Advanced

[Show additional filters](#)

Article types
Clinical Trial
Review
more ...

Text availability
Abstract available
Free full text available
Full text available

Publication dates
5 years
10 years
Custom range...

Species
Humans
Other Animals

[Clear all](#)

[Show additional filters](#)

Display Settings: Summary, 20 per page, Sorted by Recently Added

Send to:

See 2454 articles about **MTHFR** gene function
See also: **MTHFR methylenetetrahydrofolate reductase (NAD(P)H)** in the Gene database
mthfr in [Homo sapiens](#) | [Mus musculus](#) | [Rattus norvegicus](#) | [All 29 Gene records](#)

Results: 1 to 20 of 1059 << First < Prev Page 1 of 53 Next > Last >>

- ☐ [The association between MTHFR C677T polymorphism and ovarian cancer risk: a meta-analysis of 18,628 individuals.](#)
Ma C, Liu Y, Zhang W, Liu P.
Mol Biol Rep. 2013 Jan 18. [Epub ahead of print]
PMID: 23329275 [PubMed - as supplied by publisher]
[Related citations](#)
- ☐ [Short-Term Folate Supplementation in Physiological Doses Has No Effect on ESR1 and MLH1 Methylation in Colonic Mucosa of Individuals with Adenoma.](#)
Al-Ghnam Abbadi R, Emery P, Pufulete M.
J Nutrigenet Nutrigenomics. 2013 Jan 16;5(6):327-338. [Epub ahead of print]
PMID: 23328702 [PubMed - as supplied by publisher]
[Related citations](#)
- ☐ [Pharmacogenetic analysis of adjuvant FOLFOX for Korean patients with colon cancer.](#)
Lee KH, Chang HJ, Han SW, Oh DY, Im SA, Bang YJ, Kim SY, Lee KW, Kim JH, Hong YS, Kim TW, Park YS, Kang WK, Shin SJ, Ahn JB, Kang GH, Jeong SY, Park KJ, Park JG, Kim TY.

MTHFR Summary

- **chr1**
- **Size: 20,329 bp**
- **Exons: 12**
- **Potentially Functional:**
5 missense of which 3 MAF >5%
- **Previously Associated:**
3 (C677T, A1298C, A2756G)

MTHFR Example




A Deep Catalog of Human Genetic Variation

Human ▾

Search 1000 Genomes

- New Search
- Results Summary**
- Configure this page
- Add your data
- Export data
- Get VCF data
- Bookmark this page
- Share this page
- View in Ensembl



Results Summary

You searched for 'MTHFR'

Gene or Gene Product

3 entrie(s) matched your search strings.

- Gene:** [ENSG00000177000](#) Region in detail [\[1:11845780-11866977\]](#)
MTHFR - methylenetetrahydrofolate reductase (NAD(P)H) [Source:HGNC Symbol;Acc:7436]
- Variations in gene ENSG00000177000:** Variations in gene [\[1:11845780-11866977\]](#)
- Transcript:** [ENST00000376592](#) Region in detail [\[1:11845780-11863302\]](#)

Phenotype


1 entrie(s) matched your search strings.

- ENSG00000177000 PH:** [11780](#)
methylenetetrahydrofolate reductase deficiency (MTHFRD)

1000 Genomes release 16 - Oct 2014 © [EBI](#)


[About 1000 Genomes](#) | [Contact](#)

MTHFR Example



SNPs

full




102 SNPs across MTHFR → Too Many SNPs to Genotype!

Too many MTHFR SNPs

Solution: Tag SNP Selection

- SNPs are correlated (aka Linkage Disequilibrium)



Pairwise Tagging:

SNP 1

SNP 3

SNP 6

3 tags in total


Test for association:

SNP 1

SNP 3

SNP 6

Coverage: Measurement Error in TagSNPs



Want to compute coverage of all SNPs in the area, we start with this one, and look at pairwise correlation with each of the genotyped SNPs.

Genotyped SNPs

Common Measures of Coverage

- **Threshold Measures**

- e.g., 73% of SNPs in the complete set are in LD with at least one SNP in the genotyping set at $r^2 > 0.8$

- **Average Measures**

- e.g., Average maximum $r^2 = 0.84$

Coverage and Sample Size


- Sample size required for Direct Association, n
- Sample size for Indirect Association

$$n^* = n / r^2$$


- For $r^2 = 0.8$, increase is 25%
- For $r^2 = 0.5$, increase is 100%

Tag SNPs Database Resources

<http://www.hapmap.org>



<http://browser.1000genomes.org/index.html>



HapMap

- Re-sequencing to discover millions of additional SNPs; deposited to dbSNP.
- SNPs from dbSNP were genotyped
- Looked for 1 SNP every 5kb
- SNP Validation
 - Polymorphic
 - Frequency
- Haplotype and Linkage Disequilibrium Estimation
 - LD tagging SNPs
- HapMap was taken offline due to security issues & laziness (newer database)

HapMap Phase III Populations

- ASW African ancestry in Southwest USA
- CEU Utah residents with Northern and Western European ancestry from the CEPH collection
- CHB Han Chinese in Beijing, China
- CHD Chinese in Metropolitan Denver, Colorado
- GIH Gujarati Indians in Houston, Texas
- JPT Japanese in Tokyo, Japan
- LWK Luhya in Webuye, Kenya
- MEX Mexican ancestry in Los Angeles, California
- MKK Maasai in Kinyawa, Kenya
- TSI Toscani in Italia
- YRI Yoruba in Ibadan, Nigeria

1000 Genomes Project

- <http://www.internationalgenome.org/about>
- ~2-5x whole-genome sequencing coverage, some whole exome
- 2,504 individuals diverse genetic ancestries:

1000 Genomes Project

Populations

Subgroup	Supergroup											
		AFR	AMR	EAS	EUR	SAS		AFR	AMR	EAS	EUR	SAS
	ACB	96	0	0	0	0	IBS	0	0	0	107	0
	ASW	61	0	0	0	0	ITU	0	0	0	0	102
	BEB	0	0	0	0	86	JPT	0	0	104	0	0
	CDX	0	0	93	0	0	KHV	0	0	99	0	0
	CEU	0	0	0	99	0	LWK	99	0	0	0	0
	CHB	0	0	103	0	0	MSL	85	0	0	0	0
	CHS	0	0	105	0	0	MXL	0	64	0	0	0
	CLM	0	94	0	0	0	PEL	0	85	0	0	0
	ESN	99	0	0	0	0	PJL	0	0	0	0	96
	FIN	0	0	0	99	0	PUR	0	104	0	0	0
	GBR	0	0	0	91	0	STU	0	0	0	0	102
GIH	0	0	0	0	103	TSI	0	0	0	107	0	
GWD	113	0	0	0	0	YRI	108	0	0	0	0	

Haploview using 1000 Genomes Project data

(or just type in the name of the gene)

Go to the 1000 Genomes project website, and go to the 1000 Genomes Project Browser.

Enter gene name (e.g., MTHFR), to get the genomic coordinates (b37) of interest (e.g., 1:11845780-11866977) and view the region in the browser.

Click “Get VCF data” on LHS. Since you already searched for the gene (MTHFR), it will have the relevant coordinates already filled in for you.

We don’t want VCF format. Click the VCF to PED converter link on the window that pops up.

Click Next (the “Region” should have gene:coordinatesBEGIN-coordinatesEND)

Choose populations of interest – match what you are trying to target. So, for European ancestry (EUR), select CEU + FIN + GBR + IBS + TSI.

Click next, then right click on “Marker Information File” and “Linkage Pedigree File” to save the data set.

Finally, open Haploview, choose linkage format, and navigate to the pedigree information file (.ped), and then the marker information file (.info) should be auto-filled in. Click OK, and you can use Haploview!!!

Tag SNPs: HapMap & Haploview

This Column only happens because of old HapMap data (which is why I am leaving it in!) – also missing genotype rate

File Display Analysis Help

LD Plot Haplotypes **Check Markers** Tagger

#	Name	Position	Obs...	Pre...	HW...	%G...	Fam...	Me...	MAF	Rating
1	rs2184226	11781702	0.156	0.167	1.0	100.0	30	0	0.092	✓
2	rs3737967	11781715	0.094	0.101	0.263	94.4	27	1	0.054	✓
3	rs1537516	11782127	0.218	0.185	1.0	96.7	27	0	0.103	✓
4	rs1537514	11782334	0.247	0.206	0.849	98.9	29	0	0.117	✓
5	rs868014	11783713	0.0	0.0	1.0	100.0	30	0	0.0	□
6	rs4846049	11784631	0.449	0.469	1.0	98.9	29	0	0.375	✓
7	rs2274976	11785193	0.122	0.11	1.0	100.0	30	0	0.058	✓
8	rs3818762	11785269	0.448	0.428	1.0	96.7	27	0	0.31	✓
9	rs7518348	11785384	0.0	0.0	1.0	95.6	26	0	0.0	□
10	rs2274975	11785483	0.0	0.0	1.0	98.9	29	0	0.0	□
11	rs2274974	11785585	0.011	0.017	1.0	97.8	28	0	0.0090	□
12	rs2274973	11785690	0.0	0.0	1.0	100.0	30	0	0.0	□
13	rs12403856	11785777	0.0	0.0	1.0	100.0	30	0	0.0	□
14	rs6700268	11786216	0.0	0.0	1.0	98.9	29	0	0.0	□
15	rs13306556	11786376	0.233	0.206	0.849	100.0	30	0	0.117	✓
16	rs1476413	11786566	0.506	0.45	0.83	98.9	29	0	0.342	✓
17	rs17375901	11786782	0.122	0.11	1.0	100.0	30	0	0.058	✓
18	rs7535669	11787865	0.0	0.0	1.0	98.9	29	0	0.0	□
19	rs4846051	11788723	0.0	0.0	1.0	100.0	30	0	0.0	□
20	rs1801131	11788742	0.402	0.462	0.988	96.7	29	1	0.362	✓
21	rs12121543	11788937	0.404	0.399	0.552	98.9	29	0	0.275	✓
22	rs1994798	11789021	0.513	0.498	0.609	86.7	20	0	0.471	✓
23	rs2066462	11789162	0.259	0.215	0.807	94.4	25	0	0.123	✓
24	rs2066464	11789192	0.0	0.0	1.0	100.0	30	0	0.0	□
25	rs2066467	11789338	0.0	0.0	1.0	95.6	26	0	0.0	□

HW p-value cutoff: 0.0010


Min genotype %: 75

Max # mendel errors: 1 Select All

Minimum minor allele freq. 0.05

Rescore Markers

Tag SNPs: HapMap & Haploview



Tag SNPs: HapMap & Haploview

File Display Analysis Help

LD Plot Haplotypes Check Markers **Tagger**

Configuration Results

#	Name	Position	Force Incl...	Force Excl...	Capture this All...
1	rs2184226	11781702	☐	☐	☒
2	rs3737967	11781715	☐	☐	☒
3	rs1537516	11782127	☐	☐	☒
4	rs1537514	11782334	☐	☐	☒
6	rs4846049	11784631	☐	☐	☒
7	rs2274976	11785193	☒	☐	☒
8	rs3818762	11785269	☐	☐	☒
15	rs13306556	11786376	☐	☐	☒
16	rs1476413	11786566	☐	☐	☒
17	rs17375901	11786782	☐	☐	☒
20	rs1801131	11788742	☒	☐	☒
21	rs12121543	11788937	☐	☐	☒
22	rs1994798	11789021	☐	☐	☒
23	rs2066462	11789162	☐	☐	☒
28	rs6541003	11790133	☐	☐	☒
29	rs1801133	11790644	☒	☐	☒
32	rs17421462	11791113	☐	☐	☒
33	rs1572151	11791977	☐	☐	☒
34	rs17421511	11792054	☐	☐	☒
35	rs4846052	11792217	☐	☐	☒
36	rs17421560	11792590	☐	☐	☒
38	rs11121832	11794386	☐	☐	☒
39	rs2066471	11794724	☐	☐	☒
41	rs7533315	11794949	☐	☐	☒
42	rs17037390	11795109	☐	☐	☒

☒ pairwise tagging only r² threshold 0.8

☐ aggressive tagging: use 2-marker haplotypes

☐ aggressive tagging: use 2- and 3-marker haplotypes LOD threshold for multi-marker tests 3.0

Run Tagger **Reset Table**

Tag SNPs: HapMap & Haploview

File Display Analysis Help

LD Plot Haplotypes Check Markers **Tagger**

Configuration **Results**

rs2274976

rs1801131

rs1801133

rs2066462

rs2184226

rs4846052

rs17037390

rs11121832

rs2066471

rs13306561

rs12121543

rs3818762

rs1476413

rs17375901

rs9651118

Alleles captured by Current S

rs3737967

rs2274976

rs17037397

Allele	Test	r ²
rs2184226	rs2184226	1.0
rs3737967	rs2274976	0.826
rs1537516	rs2066462	1.0
rs1537514	rs2066462	1.0
rs4846049	rs1801131	0.929
rs2274976	rs2274976	1.0
rs3818762	rs3818762	1.0
rs13306556	rs2066462	0.844
rs1476413	rs1476413	1.0
rs17375901	rs17375901	1.0
rs1801131	rs1801131	1.0
rs12121543	rs12121543	1.0
rs1994798	rs4846052	0.924
rs2066462	rs2066462	1.0
rs6541003	rs4846052	1.0
rs1801133	rs1801133	1.0
rs17421462	rs2184226	1.0
rs1572151	rs2184226	1.0
rs17421511	rs2066471	1.0
rs4846052	rs4846052	1.0
rs17421560	rs2066462	0.844
rs11121832	rs11121832	1.0
rs2066471	rs2066471	1.0
rs7533315	rs11121832	0.815
rs17037390	rs17037390	1.0
rs17037396	rs2066462	1.0
rs17037397	rs2274976	1.0
rs9651118	rs9651118	1.0
rs17367504	rs17037390	1.0
rs2066470	rs2066462	1.0
rs7553194	rs2066462	0.914
rs3753582	rs2066462	0.924
rs13306561	rs13306561	1.0

Captured 33 alleles with mean r² of 0.97
 Captured 100 percent of alleles with r² > 0.8
 Using 15 SNPs in 15 tests.

Dump Tests File

Tag SNPs: HapMap Summary

- How many tag SNPs did we need, to tag all SNPs with $r^2 > 0.8$?
- In practice forced in potentially functional/previously associated SNPs
- Identified tag based on pairwise tagging
- Note: number of SNPs required varies from gene to gene and from population to population
 - Why?

Extra Slides

(if time)

Pathways

NATURE COMMUNICATIONS | DOI: 10.1038/ncomms10130

ARTICLE




Figure 3 | Ingenuity Pathway Analysis outlines potential regulatory networks around EFEMP1 and WT1. Network Analysis for EFEMP1 and WT1 was carried out using the RNA-seq FPKM > 30 gene list (see Methods section). WT1 regulates many extracellular matrix genes, including MMP2 (matrix metalloproteinase-2) and CTGF (connective tissue growth factor). EFEMP1 directly interacts with ELASTIN, a component of elastic fibres in the ECM. TIMP3 (tissue inhibitor of metalloproteinase-3), which is activated by WT1 and interacts with EFEMP1 and was found to connect between the two networks.

Pathways

- May be interested in potential joint and/or interaction effects of multiple genes in one pathway
- Many sites provide models of molecular and biochemical pathways.
- Example: BioCarta: <http://www.biocarta.com/>

Polygenic Models

- Many weak associations combine to ↑ risk?
- Score model:

$$x_j = \frac{\sum_{i=1}^m \ln(OR_i) \times SNP_{ij}}{m}$$

where

- $\ln(OR_i)$ = 'score' for SNP_i from 'discovery' sample
- SNP_{ij} = # of alleles (0,1,2) for SNP_i , person j in 'validation' sample.
- Large number of SNPs (m)
- x_j associated with disease?

Application of Model

