

Survival Analysis IV

John Kornak
April 27, 2017
john.kornak@ucsf.edu

Competing Risks

Competing Risks

- Consider a study looking at time to cardiac arrest in a high risk group comparing treatment and placebo
- But some patients die due to other causes before getting a heart attack and before being censored
- How should we treat those patients?
 - As censored?
 - As events?

Competing Risks

- Competing risks analysis mostly concerned with when non-related death can occur – potentially excluding the event of interest
- Remember conventional Kaplan-Meier – subject lost to follow up → number of subjects “at risk” drops by one
 - Assumption: independent censoring
 - Objective: extrapolate to eliminate incomplete follow up
- But what about death due to “other causes”?
- Assumption and Objective both flawed

Competing Risks

Definition: *Competing risks data* arise when multiple events can occur and follow-up can end due to occurrence of one or more of those types of events, precluding observation of at least one of the other event types

Notation:


Y : time of first event of any type

$\Delta = k$ if the k -th event type occurs first


Competing Risks

- **Elimination:** extrapolate to a scenario in which a type of event never happens [e.g., “loss to follow up” or “incomplete follow up”]
- **Accommodation:** acknowledge and allow for the competing risks – treat the competing risk as excluding the patient from having the event of interest [e.g., death]


Elimination



Accommodation



Elimination plus Accommodation



MrOS study

- Study predictors of bone fracture and low BMD
- 5993 men over 65 with average 5 year follow-up
- 531 participants fracture, 657 deaths prior to fracture, 4805 alive without fracture
- How well does baseline BMD predict fracture?

MrOS study

- How well does baseline BMD predict fracture?
- Two sources of incomplete follow-up:
 - 1) end of observation (loss to follow-up / short observation times due to staggered entry)
 - 2) death

Competing Risk Summaries

- **Cause-Specific Hazard Function** (c.f., hazard function)
- **Cummulative Incidence Function** (c.f., survival function)

Cause-Specific Hazard Function

Definition: The **cause-specific hazard function** for event type, k , $h_k(t)$, is the short-term rate at which subjects experience the onset of the k -th event among those who have not yet experienced the event of interest (e.g., fracture) or a competing event (e.g., death) prior to t .

$$\frac{\text{\# observed } k\text{-th events at time } t}{\text{\# at risk of } k\text{-th event at time } t}$$

Cause-Specific Hazard Function


observed k-th events at time t

at risk of k-th event at time t

at risk of k-th event at time t = # alive without fracture immediately prior to time t

To estimate cause-specific hazard simply set up data as ordinary survival with k-th failure as the only type – all competing causes (even death) treated as censored


Elimination



MrOS cause-specific hazard

```

use mros.dta
stset years, failure(status==1) // fracture is the event
sts graph if _t < 6.5, hazard xtitle(Years in Study)
  
```



Cause-Specific Hazard Function

Counterintuitively – **cause-specific hazard function** does not make a distinction between censoring and death...


contrast with **cumulative incidence function**...

Cumulative Incidence Function

Definition: The **cumulative incidence function** for cause k at time t, $F_k(t)$, is the proportion who have developed the k-th event prior to t

The definition above makes adjustment to “# at risk” for censoring (elimination), but not for other competing risks, i.e., for death in the MrOS dataset (accommodation)

Elimination plus Accommodation



Cumulative Incidence Function


The cumulative incidence function for the k-th event at time t , $F_k(t)$, is

$$F_k(t) = F_k(t-1) + \tilde{S}(t-1)h_k(t)$$

where $\tilde{S}(t-1)$ is the probability of being free of events at time $t-1$, i.e. just prior to time t . (In the MrOS data = chance of being alive and unfractured)

MrOS cumulative incidence function

```
clear
use mros.dta
stset years, failure(status==1)
streg , compete(status==2)
predict cif_fracture, basecif
twoway (line cif_fracture _t if _t < 6.5, sort), ///
ytitle(Cumulative Probability of Fracture) ///
xtitle(Years in Study)
```



Cox model for cause-specific hazard function

- Model cause-specific hazard function using Cox proportional hazard formulation
- Let $h_k(t)$ be the k-th cause-specific hazard, then...
- $h_k(t|x) = h_{0k}(t) \exp(\beta_{1k}x_1 + \dots + \beta_{pk}x_p)$
- Works same as original Cox model except HRs apply to the k-th cause specific hazard

MrOS Cox model for cause-specific hazard function

```
clear
use mros.dta
stset years, failure(status==1)
stcox i.bmd3 weight, nolog // bmd is defined in 3 quantiles
```

No. of subjects = 5993 Number of obs = 5993
 No. of failures = 531
 Time at risk = 30483.46339
 Log likelihood = -4442.2904 LR chi2(3) = 121.22
 Prob > chi2 = 0.0000

_t	Haz. Ratio	Std. Err.	z	P> z	[95% Conf. Interval]
bmd3					
2	.4193745	.0447953	-8.14	0.000	.3401585 .5170384
3	.3290229	.0396476	-9.23	0.000	.2598098 .4166743
weight	1.004146	.00362	1.15	0.251	.997076 1.011266

Fine-Gray model for cumulative incidence function

- What if BMD is correlated with factors that make death more likely?
- Low BMD might lead to fewer fractures after 2 years because men with low BMD are more likely to die before fracture.
- Solution: compare cumulative incidence function rather than cause-specific hazard.

Fine-Gray model for cumulative incidence function

- Model cumulative incidence function, adapting Cox proportional hazard formulation
- Let $f_k(t)$ be the k-th "hazard" where the cohort members who succumb to the competing risk are retained in the denominator of their rate (c.f., death)
- $f_k(t|x) = f_{0k}(t) \exp(\beta_{1k}x_1 + \dots + \beta_{pk}x_p)$
- Note: Maintaining deaths in the risk set acknowledges someone lost to competing risk will not develop event of interest so no extrapolation required

MrOS Fine-Gray model for cumulative incidence function

```
clear
use mros.dta
stset years, failure(status==1)
stcrreg i.bmd3 weight, compete(status==2) // bmd in 3 quantiles
```

Competing-risks regression No. of obs = 5993
 No. of subjects = 5993
 Failure event : status == 1 No. failed = 531
 Competing event: status == 2 No. competing = 657
 No. censored = 4805

Log pseudolikelihood = -4472.9261 Wald chi2(3) = 119.64
 Prob > chi2 = 0.0000

_t	SHR	Robust Std. Err.	z	P> z	[95% Conf. Interval]
bmd3					
2	.4219663	.0443983	-8.20	0.000	.3433337 .5186079
3	.3369128	.03982	-9.20	0.000	.2672472 .4247387
weight	1.004669	.0036759	1.27	0.203	.9974896 1.011899

MrOS model comparisons

Cause-specific Cox model

_t	Haz. Ratio	Std. Err.	z	P> z	[95% Conf. Interval]	
bmd3						
2	.4193745	.0447953	-8.14	0.000	.3401585	.5170384
3	.3290229	.0396476	-9.23	0.000	.2598098	.4166743
weight	1.004146	.00362	1.15	0.251	.997076	1.011266

Fine-Gray model

_t	SHR	Robust Std. Err.	z	P> z	[95% Conf. Interval]	
bmd3						
2	.4219663	.0443983	-8.20	0.000	.3433337	.5186079
3	.3369128	.03982	-9.20	0.000	.2672472	.4247387
weight	1.004669	.0036759	1.27	0.203	.9974896	1.011899

MrOS model comparisons

- MrOS cause-specific hazards and Fine-Gray model results are very similar
- Makes sense: Methods differ based on whether people who die are retained in the risk set – but the risk of death is not large.
- Neither approach is incorrect; all depends on desired interpretation

Competing Risks Summary

- “Deaths” occur due to other causes before event of interest
- Elimination vs. Accommodation
- Cause-specific hazard; cause-specific Cox model (stcox); elimination only
- Cumulative incidence function; Fine-Gray model (stcrreg); elimination for censoring and accommodation for competing death (by other causes)

Finally...

- Dr. McCulloch's first lecture on Repeated Measures coming Tuesday
- Check syllabus – all solutions (labs + homeworks) for my lectures should be posted by weekend – let me know if not
- Final project assignments will be in place today