

Common Biostatistical Problems And the Best Practices That Prevent Them

Biostatistics 209
April 21, 2020

Chiung-Yu Huang

(originally developed by Peter Bacchetti)

- HW#2 due today


Goal: Provide conceptual and practical dos, don'ts, and guiding principles that help in

- Choosing the most meaningful analyses
- Understanding what results of statistical analyses imply for the issues being studied
- Producing clear and fair presentation and interpretation of results

There may be exceptions

Please let me know about additions or disagreements during lecture, or later (ChiungYu.Huang@ucsf.edu)


If you find this lecture interesting ...
Optional reference text -- short and very readable
(See especially chapters 12, 15, 31, 33)



Amazon.com link: http://www.amazon.com/p-value-Stories-Actually-Understand-Statistics/dp/0321629302/ref=sr_1_1?ie=UTF8&s=books&qid=1270360017&sr=8-1

In 2015, a psychology journal announced that it will no longer publish papers containing P values because the statistics were too often used to support lower-quality research.

In response to this, the American Statistical Association (ASA) issued the statement on p-values:



The American Statistician



ISSN: 0003-1305 (Print) 1537-2731 (Online) Journal homepage: <http://amstat.tandfonline.com/loi/utas20>

The ASA's Statement on p-Values: Context, Process, and Purpose

Ronald L. Wasserstein & Nicole A. Lazar

To cite this article: Ronald L. Wasserstein & Nicole A. Lazar (2016) The ASA's Statement on p-Values: Context, Process, and Purpose, The American Statistician, 70:2, 129-133, DOI: [10.1080/00031305.2016.1154108](https://doi.org/10.1080/00031305.2016.1154108)

ASA's Statement on p-Values

“The p-value was never intended to be a substitute for scientific reasoning”

“Well-reasoned statistical arguments contain much more than the value of a single number and whether that number exceeds an arbitrary threshold.”

In 2019, the American Statistician published a special issue “Moving to a World Beyond $p < 0.05$ ” explaining how to use p-values responsibly



The American Statistician



ISSN: 0003-1305 (Print) 1537-2731 (Online) Journal homepage: <https://tandfonline.com/loi/utas20>

Moving to a World Beyond “ $p < 0.05$ ”

Ronald L. Wasserstein, Allen L. Schirm & Nicole A. Lazar

To cite this article: Ronald L. Wasserstein, Allen L. Schirm & Nicole A. Lazar (2019) Moving to a World Beyond “ $p < 0.05$ ”, The American Statistician, 73:sup1, 1-19, DOI: [10.1080/00031305.2019.1583913](https://doi.org/10.1080/00031305.2019.1583913)

To link to this article: <https://doi.org/10.1080/00031305.2019.1583913>

“We conclude... that it is time to stop using the term ‘statistically significant’ entirely. Nor should variants such as ‘significantly different,’ ‘ $p < 0.05$,’ and ‘nonsignificant’ survive, whether expressed in words, by asterisks in a table, or in some other way.”

“Where p-values are used, they should be reported as continuous quantities”

“Accept uncertainty. Be thoughtful, open, and modest”

What is a p-Value?

A p-value is the probability under a specified statistical model that a statistical summary of the data would be equal to or more extreme than its observed value.

E.g. Under the null hypothesis of no difference between groups, the standardized sample mean difference follows a standard normal distribution when N is sufficiently large. If we observed a standardized difference of 1, the two-sided p-value is $P(|Z| \geq 1) = 0.32$.

P-values – TRUE or FALSE?

The p-value tells you whether an observed difference, effect, or association is real or not.

P-values – TRUE or FALSE?

If the result is not statistically significant, that proves there is no difference.

P-values – TRUE or FALSE?

If the result is not statistically significant, you have to conclude that there is no difference.

Problem 1. P-values for establishing negative conclusions

The P-value Fallacy:

The p-value tells you whether an observed difference, effect, or association is real or not.

If the result is not statistically significant, that proves there is no difference.

If the result is not statistically significant, you have to conclude that there is no difference.

-- ALL FALSE

How about:

$p > 0.05$ + Power Calculation = No effect

How about:

$p > 0.05$ + Power Calculation = No effect

Still no good!

Reasoning via p-values and power is convoluted and unreliable.

Power calculations are usually inaccurate. They rely heavily on assumptions that aren't known in advance. Inaccuracy is theoretically inevitable and empirically verified.

A study of RCTs in 4 top medical journals found more than half used assumed SD's off by enough to produce >2-fold differences in sample size.

CONSORT (Consolidated Standards of Reporting Trials) guidelines for reporting RCTs warn against this practice: “There is little merit in a post hoc calculation of statistical power using the results of a trial”.

A simpler and better alternative:
Confidence intervals show simply and directly what possibilities are reasonably consistent with the observed data.

Additional references for not using power argument:

1958, D.R. Cox: “Power . . . is quite irrelevant in the actual analysis of data.”

Goodman SN, Berlin JA. The use of predicted confidence intervals when planning experiments and the misuse of power when interpreting results. *Ann Intern Med* 1994; **121**:200-6.

Hoening JM, Heisey DM. The abuse of power: the pervasive fallacy of power calculations for data analysis. *American Statistician*. 2001;**55**:19-34.

Bacchetti P. Current sample size conventions: flaws, harms, and alternatives. *BMC Medicine*, **8**:17, 2010.

How about:

$p > 0.05$ + Large N = No effect

$p > 0.05$ + Huge Expense = No effect

$p > 0.05$ + Massive Disappointment = No Effect

How about:

$p > 0.05$ + Large N = No effect

$p > 0.05$ + Huge Expense = No effect

$p > 0.05$ + Massive Disappointment = No Effect

Temptation to make the results sound stronger --
Not if contradicted by the CI's!

Confidence intervals show simply and directly what values are reasonably consistent with the observed data.

Example: Treatment of an acute infection

Treatment A: 16 deaths in 100

Treatment B: 8 deaths in 100

Odds ratio:

2.2, 95% CI 0.83 to 6.2, $p=0.13$

Estimated risk difference:

8.0%, 95% CI -0.9% to 16.9%

Example: Treatment of an acute infection

Treatment A: 16 deaths in 100

Treatment B: 8 deaths in 100

Odds ratio:

2.2, 95% CI 0.83 to 6.2, $p=0.13$

Risk difference:

8.0%, 95% CI -0.9% to 16.9%

“No difference in death rates”

“No **significant** difference in death rates”

“No **statistical** difference in death rates”

“No **statistically significant** difference in death rates”

-- but the difference was “significant” to the estimated 8 additional people who died with treatment A.

Example: Treatment of an acute infection

Treatment A: 16 deaths in 100

Treatment B: 8 deaths in 100

Odds ratio: 2.2, 95% CI 0.83 to 6.2, $p=0.13$

Risk difference: 8.0%, 95% CI -0.9% to 16.9%

A more sensible interpretation:

“Our study suggests an important benefit of Treatment B, but this did not reach statistical significance.”

NEJM, **354**: 1796-1806, 2006.

“Supplementation with vitamins C and E during pregnancy *does not reduce* the risk of preeclampsia in nulliparous women, the risk of intrauterine growth restriction, or the risk of death or other serious outcomes in their infants.”

Preeclampsia: RR 1.20 (0.82 – 1.75)

Growth restriction: RR 0.87 (0.66 – 1.16)

Serious outcomes: RR 0.79 (0.61 – 1.02)

Women's Health Initiative study on fat consumption and breast cancer

HEALTH

THE NEW FIGHT OVER FAT

BY JERRY ADLER

IF YOU WERE WONDERING what to make of the definitive eight-year study on dietary fat by the Women's Health Initiative released last week, you're not alone. Even some leading researchers were having trouble figuring out what to say about the study's major conclusion: that a low-fat diet did not significantly reduce disease among nearly 20,000 postmenopausal women, compared with a control group who ate what they wanted.

Was Ross L. Prentice of the Fred Hutchinson Cancer Research Center, one of the authors of the study, sounding slightly defensive when he proclaimed that "women can be confident that cutting back on fat... certainly won't hurt when it comes to maintaining a healthy lifestyle"? (Emphasis added.) Did the food industry waste the billions it spent inventing fat-free cookies?

Well, maybe. The problem, says Dr. Marcia Stefanick of Stanford, who heads the steering committee of the WHI, is that the study was designed back in the early 1990s to test an idea that most researchers were already starting to abandon: that the key to health is the total amount of fat in your diet. Instead, most nutritionists now emphasize controlling calories and eating healthy fats—olive and other unsaturated vegetable oils—while avoiding the bad kinds. So it was no great surprise when The Journal of the American Medical Association reported that researchers had

found minor reductions, or none at all, in breast or colon cancer or heart disease among women who cut their fat intake on average to less than 29 percent of total calories. (The control group ate a typical American diet with 35 to 38 percent fat.) Those results "are very consistent with what we've seen" in research over the past decade, says Dr. Walter Willett, the prominent Harvard nutritionist, who calls the craze for low-fat everything a "distraction" from good dietary advice.

And that advice—for both women and men—is just what you've been hearing for

Even diets with only 29% of calories coming from fat didn't reduce the risk of disease.



the past decade: to avoid trans fats (the partially hydrogenated vegetable oils found in processed foods) and restrict saturated fats from meat and dairy products, while consuming a healthy balance of vegetables, fruits and whole grains. "People should stop thinking low fat is the same as healthy," says Stefanick. "The food industry did a great job of selling that, and people believed them." The other advice from nutritionists hasn't changed, either: to exercise and control total calories to avoid obesity.

Exercise is important even apart from its effect on weight: it regulates glucose metabolism (lowering the risk of diabetes) and improves bowel function (which may cut the risk of colon cancer). Obesity appears to cause hormonal changes implicated in breast cancer in postmenopausal women, notes Dr. Michael Thun of the American Cancer Society. In the study, the women who ate a lower-fat diet didn't lose weight, but neither did they gain—a fact that gives small comfort to either side in the great struggle between the authors of low-fat and low-carb diet books.

Even after this definitive study, though, most nutritionists (except for those in the Atkins ultra-low-carb camp) still think there's a benefit to limiting fat consumption. Buried in the larger story of the study was the intriguing statistic that

Invasive Breast Cancer
HR 0.91 (0.83-1.01),
p=0.07

Breast Cancer Mortality
HR 0.77 (0.48-1.22)

From *JAMA* abstract:
"a low-fat dietary pattern did not result in a statistically significant reduction in invasive breast cancer risk"

Read Dr. Dean Ornish's new column on dieting, nutrition and health.

Check out the best Web sites for learning about Black History Month.

Read more about the Hazelden drug center at Newsweek.com on MSNBC.

PHOTO ILLUSTRATION BY NEWSWEEK. PHOTOS (FROM LEFT): HEMERA TECHNOLOGIES—ALAMY, ELIZABETH WATTS—JUPITER IMAGES

FEBRUARY 20, 2006 NEWSWEEK 69

Best Practice 1. Provide estimates—with confidence intervals—that directly address the issues of interest.

Often followed (but then ignored when interpreting)

BP2. Ensure that major conclusions reflect the estimates and the uncertainty around them.

BP2a. Never interpret large p-values as establishing negative conclusions.

The estimates and CI's should contribute to the interpretation, not just the p-value:

- The estimate is the value most supported by the data
- The confidence interval includes values that are not too incompatible with the data
- The study provides strong evidence against values outside the CI

Example of a well supported strong negative result:

NEJM, **354**: 1889-1900, 2006

Conclusion: “When treated with phototherapy or exchange transfusion, total serum bilirubin levels in the range included in this study were not associated with adverse neurodevelopmental outcomes in infants born at or near term.”

Support: “on most tests, 95 percent confidence intervals excluded a 3-point (0.2 SD) decrease in adjusted scores in the hyperbilirubinemia group.”

What if results are less conclusive?

Growth restriction: RR 0.87 (0.66 – 1.16)

Serious outcomes: RR 0.79 (0.61 – 1.02)

“Our results suggest that Vitamin C and E supplementation may substantially reduce the risk of growth restriction and the risk of death or other serious outcomes in the infant, but confidence intervals were too wide to rule out the possibility of no effect.”

But then the paper probably won't end up in NEJM!

The “elephant in the room” when it comes to conflict of interest:

- We are all under pressure to make our papers seem as interesting as possible.
- Under-recognized conflict of interest

The p-value fallacy can help make “negative” studies seem more conclusive and interesting.

Be vigilant (and be honest)!

BP3. Discuss the implications of your findings for what may be true in general. Do not focus on “statistical significance” as if it were an end in itself.

WHI conclusion:

“a low-fat dietary pattern **did not result in a statistically significant reduction** in invasive breast cancer risk ... However, the nonsignificant trends ... indicate that longer, planned, nonintervention follow-up may yield a **more definitive comparison.**”

Newsweek followup article:

“The conclusion of the breast-cancer study—that a low-fat diet **did not lower** risk—was fairly nuanced. It suggested that if the women were followed for a longer time, there might be **more of an effect.**”

Easy to slip into relying on “ $p>$ ” reasoning

- Yes or No reasoning more natural
- Focus on p-values engrained in research culture
- Real level of uncertainty often inconveniently large, which can make results seem less interesting

Be vigilant

- Double-check all negative interpretations
- Examine estimates, confidence intervals

How to check negative interpretations:

Perform searches for words “no” and “not”


Check each negative interpretation found

- Is there an estimate and CI supporting this?
- What if the estimate were exactly right?
- What if the upper confidence bound were true?
- What if the lower confidence bound were true?


Additional searches: “failed”, “lack”, “absence”, “disappeared”, “only”, “rather”, “neither”, “none”

Drawing conclusions from CIs


A somewhat idealized situation where there is consensus on what is clinically important




We found strong evidence against any substantial harm or benefit.




Estimate suggests substantial benefit
CI suggests may be no effect (not statistically significant)



Strong evidence of benefit (statistically significant)
Substantial benefit appears likely, but CI too wide to
rule out clinically unimportant benefit



Strong evidence of substantial clinical benefit



No conclusions possible due to very wide CI

Also see online resource

<http://ctspedia.org/do/view/CTSpedia/ResultsInterpretation>

**CTSPEDIA**

You are here: [CTSpedia](#) > [CTSpedia Web](#) > [ContentInterest](#) > [ResultsInterpretation](#) > [InterpretationText](#)
(07 Oct 2014, [PeterBacchetti](#))

Tags:
[create new tag](#), [view all tags](#), [tagging instructions](#)

[Edit](#) [Attach](#)

Interpretation of Coefficients

Directions

This page will help you write an interpretation of an estimate that considers both its substantive and statistical significance.

Interpretation requires assessing whether the effect would be large enough to be important if it turned out to be exactly right. "Important" can mean important for patients, for public health, or for scientific understanding. Similar assessments are also needed for the confidence bounds. Use the pull-down menus to provide these assessments.

When you have filled in all the values, suggested text appears below the table. You can change the values and the text will be rewritten appropriately. You will likely need to customize the text to match your particular situation.

Fill in Here

Was your estimate statistically significant?

	Value	Importance
Estimate		Please select interpretation
Lower Confidence Limit		Please select interpretation
Upper Confidence Limit		Please select interpretation

CTSpedia
[Log In or Register](#)

Webs on CTSpedia

- [CTSpedia](#)
- [Main](#)
- [ResearchEthics](#)
- [Sandbox](#)
- [System](#)

[Recent Changes](#)
[Help](#)

Example from a typical collaboration:

First draft text:

“There were no statistically significant effects of DHEA on lean body mass, fat mass or bone density.”

Final wording:

“Estimated effects of DHEA on lean body mass, fat mass, and bone density were small, but the confidence intervals around them were too wide to rule out effects large enough to be important.”

Are large p-values good for anything?

“Due diligence” situations

Checking for possible model assumption violations
when little suspicion

Just need to state that you checked and nothing jumped out; don't need to prove that no violation was possible:

“We note that the confidence intervals were not narrow enough to rule out potentially important interactions, but in the absence of strong evidence for such interactions we focus on the simpler models without them.”

Problem 2. Misleading and vague phrasing

We failed to detect ...

Our results do not support ...

We found no evidence for ...

Our data did not confirm ...

“There is no scientific evidence that BSE [Mad Cow Disease] can be transmitted to humans or that eating beef causes it in humans.”

-- British Prime Minister John Major, 1995

Preliminary investigations conducted by the Chinese authorities have found no clear evidence of human-to-human transmission of the novel coronavirus (2019-nCoV) identified in Wuhan, China

-- WHO, Jan 14, 2020

BP4. State what you did find or learn, not what you didn't.

This prevents deception, but also can make statements clearer and stronger.

Oddly, investigators often understate their conclusions using weak phrasing.

FRAM, nationwide study of fat abnormalities in persons with HIV

Peripheral fat loss association with central fat gain, OR: 0.71, CI: 0.47 to 1.06, $P = 0.10$.

First draft: “our results do not support the existence of a single syndrome with reciprocal findings.”

Final: “We found evidence against any reciprocal increase in VAT in HIV-infected persons with peripheral lipoatrophy”

Safety of cannabinoids in persons with treated HIV

Marijuana effect on $\log_{10}$ VL: -0.06 (-0.26 to 0.13)
Dronabinol: -0.07 (-0.24 to 0.06)

First draft: “Overall there was no evidence that cannabinoids increased HIV RNA levels over the 21-day study period.”

Final: “This study provides evidence that short-term use of cannabinoids, either oral or smoked, does not substantially elevate viral load in individuals with HIV infection.”

Problem 3. Speculation about low power

Sometimes investigators want to argue that their hoped-for results could still be possible

“There were departures from the design assumptions that likely reduced study power.

...

“If the WHI design assumptions are revised to take into account these departures [less dietary fat reduction], projections are that breast cancer incidence in the intervention group would be 8% to 9% lower than in the comparison group [and] the trial would be somewhat underpowered (projected power of approximately 60%) to detect a statistically significant difference, which is consistent with the observed results.”

What are they trying to say?

There might be a 9% reduction in risk. We could have missed it because power was only 60%.

But estimated $HR = 0.91$, so of course a 9% reduction is possible. It's what they actually saw!

BP2. Ensure that major conclusions reflect the estimates and the uncertainty around them.

Problem 4. Exclusive reliance on intent-to-treat analysis

‘Negative’ study of vitamin E in diabetics (*JAMA* 2005)

“To reduce bias, we included continuing followup from those who declined active participation in the study extension and stopped taking the study medication.”

But ITT produces underestimates of actual biological effects: it is biased toward no effect.

WHI:

Estimate of effect if adherent to low-fat diet:

Breast cancer HR 0.85 (0.71 – 1.02)

Use of more stringent adherence definition “leads to even smaller HR estimates and to 95% CIs that exclude 1.”

BP5. Learn as much as you can from your data.

BP5a. Also consider secondary per-protocol analyses, especially if:

- Interest in biological issues
- Double-blinded treatment

BP5b. Consider advanced methods to estimate causal effects.

Problem 5. Reliance on omnibus tests

Problem 6. Overuse of multiple comparisons adjustments

Omnibus tests (like ANOVA)

- check for any one or more of a large number of possible departures from a global null hypothesis (nothing is happening anywhere)
- inherently focused only on p-values (Problem 1)
- diffuse, so weaker for specific issues

Multiple comparisons adjustments

- each result detracts from the other
- with no consideration of how well the different results fit together conceptually or scientifically

Investigator's panicked inquiry:

Animal experiment that included

- a condition that just confirms that the experiment was done correctly
- some places where different conditions should be similar
- some conditions that should differ

Saw expected results in pairwise comparisons, but
“ANOVA says that there is nothing happening”

Reviewer's comment on a study examining effects of 4 different administration routes

“Repeated measures analysis of variance should be completed. Only if the time-by-treatment interaction is significant, should time-specific comparisons be made. Then multiple comparison procedures, such as Tukey's test, should be used rather than repeated t tests.”

This would treat $p > 0.05$ on the unfocused omnibus test of time-by-treatment interaction as a reliable indicator that no important differences are present—**Problem 1**.

Study of biology of morphine addiction:

Very complex design involving:

- two different receptors
- antagonists
- different brain regions with and w/o certain receptor
- systemic vs local administration

Results of many pairwise comparisons fit a biologically coherent pattern.

Reviewer: “The statistical analyses are naïve. The authors compute what appear to be literally dozens of t-tests without any adjustment to the alpha level --- indeed the probability of obtaining false positives grows with the number of such tests computed. The authors should have conducted ANOVAs followed by the appropriate post-hoc tests. Their decision to simply compute t-tests on all possible combinations of means is statistically unacceptable.”

But the probability of obtaining multiple positive results exactly where expected and negative results exactly where expected does not grow; it becomes vanishingly small.

BP6. Base interpretations on a synthesis of statistical results with scientific considerations.

BP6a. Rely on scientific considerations to guard against overinterpretation of isolated findings with $p < 0.05$. (This is usually preferable to formal multiple comparisons adjustment.)

BP6b. Acknowledge the desirability of independent replication, particularly for unexpected findings.

BP7. Choose accuracy over conservatism whenever possible.

Problem 7. Entangled outcomes and predictors

Total time on treatment as a (fixed) predictor of survival time

Can only be treated if alive
(survival time = treated time + untreated time)

Died after 2 days $\rightarrow$ max of 2 days treatment
Treated for 5 years $\rightarrow$ min of 5 years survival

Meaningless association

BP 8: Either

- 1) ensure that outcome is not part of the definition of a predictor, and vice versa, or
- 2) be very careful and clear with interpretation

Fixed (e.g. baseline) predictors should be defined without using any information from after the start of followup

Use time-dependent covariates, defined only using measurements up to present

Technical problems

Unchecked assumptions

- Always list assumptions, describe checks and results thereof

Ignored dependence and clustering

- e.g., unpaired analysis on paired (before/after) data
- Dr. McCulloch's lectures deal with more complex levels of clustering and repeated measures

Unclear details for time-to-event

- operational definitions – start time, event occurrence, censoring time
- summarize follow up among those censored
- summarize early loss to follow up separately
- how were events ascertained?

Poor summaries

- e.g., mean $\pm$ SD for skewed data – better median/IQR
- SD's or SE's? SE's (or better CI's) for precision of estimates, whereas SD's for population variability

Showing inadequate or excessive precision

- too little $\Rightarrow$ reader wonders how big exactly
- too much $\Rightarrow$ false impression of precision (naïve)
- p-values 2 significant digits, HR's 2 decimals if < 2.0 and one if ≥ 2.0
- other values – means, SDs etc., balance scientific relevance, clarity and realistic accuracy
- not $p=\text{NS}$ or $p>0.1$ – encourages p-value fallacy
- do not show χ^2 , F statistics – nothing over p-value

Poorly scaled predictors

- e.g., don't look at age effects per day... in particular HR's difficult to convert in the reader's head.

Terms likely to be misread

- especially “significant” – use “statistically significant”, or if implying more general definition use “important”, “substantial”, or (best) “clinically significant” – if journal insists on “significant” for “statistically significant” try to avoid completely.
- use “relative hazard” or “hazard ratio” for proportional hazards models – NOT “relative risk” – latter used for relative risk modeling of binary outcomes.
- Use “Mann-Whitney” or “Wilcoxon rank-sum” not “Wilcoxon” to avoid confusion with “Wilcoxon signed-rank”.

Missing Data

- can cause bias -- depends on why missing
- e.g., those with the riskiest behavior may be more inclined to skip questions about risk factors.

Prevention is best, otherwise

- clearly disclose how much
- give reasons when known
- assess differences in other variables of non-responders
- perform sensitivity analyses – compare different assumptions for the missing data. Do conclusions change?

- consider advanced methods, e.g., Stata can perform *multiple imputation* (see the *mi* command and <http://www.ats.ucla.edu/stat/stata/Library/ice.htm>). You will likely need help from a statistician to use such approaches.
- when comparing single and multiple predictor model results, use same set of observations.

Homework

Examine the two assigned papers

Look for:

- use of best practices, other strengths
- problems
- missed opportunities for using best practices

Think about what would have been better and the practical or scientific consequences

We will discuss these on Thursday

Written Homework

Due **before class THIS Thursday 4/23**

Choose one paper and

describe occurrence of one of the problems. Quote text or pinpoint specific results or passages to clearly identify what you discuss. What should have been done to avoid the problem?

Heisler M, Faul JD, Hayward RA, Langa KM, Blaum C, Weir D. Mechanisms for Racial and Ethnic Disparities in Glycemic Control in Middle-aged and Older Americans in the Health and Retirement Study. *Arch Intern Med* 2007; **167**:1853-1860.

Homsy J, Bunnell R, Moore D, King R, Malamba S, et al. Reproductive Intentions and Outcomes among Women on Antiretroviral Therapy in Rural Uganda: A Prospective Cohort Study. *PLoS ONE* 2009; **4**(1): e4149.

Summary of Problems

- Problem 1.** P-values for establishing negative conclusions
- Problem 2.** Misleading and vague phrasing
- Problem 3.** Speculation about low power
- Problem 4.** Exclusive reliance on intent-to-treat analysis
- Problem 5.** Reliance on omnibus tests
- Problem 6.** Overuse of multiple comparisons adjustments
- Problem 7.** Entangled outcomes and predictors

Summary of Biostatistical Best Practices

BP1. Provide estimates—with confidence intervals—that directly address the issues of interest.

BP2. Ensure that major conclusions reflect the estimates and the uncertainty around them.

BP2a. Never interpret large p-values as establishing negative conclusions.

BP3. Discuss the implications of your findings for what may be true in general. Do not focus on “statistical significance” as if it were an end in itself.

BP4. State what you did find or learn, not what you didn’t.

Summary of Biostatistical Best Practices

BP5. Learn as much as you can from your data.

BP5a. Also consider per-protocol analyses, especially if:

- Interest in biological issues
- Double-blinded treatment

BP6. Base interpretations on a synthesis of statistical results with scientific considerations.

BP6a. Rely on scientific considerations to guard against overinterpretation of findings with $p < 0.05$.

BP6b. Acknowledge the desirability of independent replication, particularly for unexpected findings.

BP7. Choose accuracy over conservatism whenever possible.

BP8. Ensure outcome is not part of the definition of a predictor, or (if you must) be very careful and clear with interpretation.

Specific exercise for written projects:

Perform searches for words “no” and “not”

Check each sentence found

- Is there an estimate and CI supporting this?
- What if the point estimate were exactly right?
- What if the upper confidence bound were true?
- What if the lower confidence bound were true?

Additional searches: “failed”, “lack”, “absence”,
“disappeared”, “only”, “rather”, “neither”, “none”

Also for your written projects

Try to avoid the other problems and follow the best practices

(Or be clear on why your case is an exception)

Take advantage of the faculty help that is available