

PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

Improving Situation Awareness to Reduce Unrecognized Clinical Deterioration and Serious Safety Events

Patrick W. Brady, Stephen Muething, Uma Kotagal, Marshall Ashby, Regan Gallagher, Dawn Hall, Marty Goodfriend, Christine White, Tracey M. Bracke, Victoria DeCastro, Maria Geiser, Jodi Simon, Karen M. Tucker, Jason Olivea, Patrick H. Conway and Derek S. Wheeler

Pediatrics 2013;131:e298; originally published online December 10, 2012;

DOI: 10.1542/peds.2012-1364

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://pediatrics.aappublications.org/content/131/1/e298.full.html>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2013 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Improving Situation Awareness to Reduce Unrecognized Clinical Deterioration and Serious Safety Events

abstract

BACKGROUND AND OBJECTIVE: Failure to recognize and treat clinical deterioration remains a source of serious preventable harm for hospitalized patients. We designed a system to identify, mitigate, and escalate patient risk by using principles of high-reliability organizations. We hypothesized that our novel care system would decrease transfers determined to be unrecognized situation awareness failures events (UNSAFE). These were defined as any transfer from an acute care floor to an ICU where the patient received intubation, inotropes, or ≥ 3 fluid boluses in first hour after arrival or before transfer.

METHODS: The setting for our observational time series study was a quaternary care children's hospital. Before initiating tests of change, 2 investigators reviewed recent serious safety events (SSEs) and floor-to-ICU transfers. Collectively, 5 risk factors were associated with each event: family concerns, high-risk therapies, presence of an elevated early warning score, watcher/clinician gut feeling, and communication concerns. Using the model for improvement, an intervention was developed and tested to reliably and proactively identify patient risk and mitigate that risk through unit-based huddles. A 3-times daily inpatient huddle was added to ensure risks were escalated and addressed. Later, a "robust" and explicit plan for at-risk patients was developed and spread.

RESULTS: The rate of UNSAFE transfers per 10 000 non-ICU inpatient days was significantly reduced from 4.4 to 2.4 over the study period. The days between inpatient SSEs also increased significantly.

CONCLUSIONS: A reliable system to identify, mitigate, and escalate risk was associated with a near 50% reduction in UNSAFE transfers and SSEs. *Pediatrics* 2013;131:e298–e308

AUTHORS: Patrick W. Brady, MD, MSc,^{a,b} Stephen Muething, MD,^{a,b} Uma Kotagal, MBBS, MSc,^b Marshall Ashby, MHA, MBA,^b Regan Gallagher, MSN,^e Dawn Hall, BSN, MHA,^e Marty Goodfriend, BSN, MEd,^{d,e} Christine White, MD, MAT,^a Tracey M. Bracke, MS,^b Victoria DeCastro, MSN, MBA,^e Maria Geiser, BSN, Jodi Simon, MHA,^f Karen M. Tucker, MSN, RN,^e Jason Olivea, MS, MPA,^b Patrick H. Conway, MD, MSc,^{a,g} and Derek S. Wheeler, MD^{b,c}

Divisions of ^aHospital Medicine, ^cCritical Care Medicine, Department of Pediatrics, and ^dFamily Relations, ^bThe James M. Anderson Center for Health Systems Excellence, ^eDepartment of Patient Services, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio; ^fDivision of Quality Services, Akron Children's Hospital, Akron, Ohio; and ^gCenters for Medicare and Medicaid Services, Office of Clinical Standards and Quality, Baltimore, Maryland

KEY WORDS

patient safety, situation awareness, rapid response systems, clinical deterioration, quality improvement, high-reliability organizations, hospital medicine

ABBREVIATIONS

ACA—apparent cause analysis
EHR—electronic health record
HRO—high-reliability organization
MPS—manager of patient services
MRT—medical response team
PEWS—pediatric early warning score
RRT—rapid response team
SA—situation awareness
SOD—safety officer of the day
SSE—serious safety event
UNSAFE—unrecognized situation awareness failure event

(Continued on last page)

Rapid response teams (RRTs) are designed to identify and respond to events in the hours prearrest.^{1–13} Although interventions and contexts have varied substantially, these teams have demonstrated decreased codes outside the ICU and hospital-wide mortality in several studies.^{1–4,7,9,10,12,13} Variation in the effectiveness of RRTs may be due to insufficient processes around monitoring and risk identification.¹⁴ Potentially preventable morbidity and mortality from unrecognized deterioration remain, often due to ineffective clinical monitoring that we believe represents poor situation awareness (SA).¹⁴ SA (ie, “knowing what is going on”) exists at 3 levels and is defined as “the perception of elements in the environment within a volume of time and space, the comprehension of their meaning, and the projection of their status in the near future.”^{15–17} We believe improved SA drives better recognition of early deterioration and is essential in efforts to reduce “failure to rescue” from codes outside of the ICU, an event associated with a 50% to 67% mortality.^{18,19}

High-reliability organizations (HROs) (eg, commercial aviation, nuclear power, and wilderness firefighting) deal with constant and catastrophic risk yet maintain exemplary safety records.²⁰ Our institution began a journey to become an HRO with the Agency for Healthcare Research and Quality HRO Learning Network in September 2005.²¹ Learnings from this network fueled improvement work and introduced the concept of SA. Our organization has targeted serious safety event (SSE) reduction as a strategic improvement goal since 2006. Our efforts to reduce SSEs, defined as severe harm or death after variation from expected practice, have resulted in significant and sustained reduction.²² Before SA work, SSEs among inpatients had not decreased. Poor SA was a common

etiology. To achieve our aim and facilitate rapid learning, our project team defined a precursor outcome measure that we believed would capture events that both represented SA failures and occurred more commonly than inpatient SSEs in our center.²³ We prospectively defined unrecognized situation awareness failures events (UNSAFE) as the transfer of patients from the acute care floor to the ICU where the patient received tracheal intubation, initiation of vasoactive medications for hemodynamic support, or ≥ 3 fluid boluses in the first 60 minutes of ICU care or before arrival in the ICU. We believed these events represented potentially delayed transfers that are precursors to codes outside the ICU or SSEs. With focused improvement work beginning in November 2009, we aimed to decrease UNSAFE transfers by 50% by June 30, 2011. We hypothesized that a system of care that proactively identified, mitigated, and escalated risk would improve SA and decrease UNSAFE transfers and SSEs.

METHODS

Setting

Cincinnati Children's Hospital Medical Center is a 523-bed academic, quaternary-care, free-standing children's hospital. Resident teaching teams care for the majority of hospitalized patients. Less commonly, direct hospitalist and nurse-practitioner models are used. Our RRT (called a medical response team [MRT]) has been in place since 2006 with defined activation criteria.² A modified version of the Monaghan pediatric early warning score (PEWS) was tested and spread across the hospital in 2007.²⁴

Human Subjects Protection

Our study was reviewed by the institutional review board and deemed exempt systems improvement. Individual

patient care was discussed among clinicians in the course of identification and mitigation of risk. Medical record review was performed by the lead investigator by using a secure password-protected internal database and our hospital's electronic health record (EHR).

Event Review

Two investigators (Dr Brady and Ms Goodfriend) reviewed 20 consecutive SSEs and 80 consecutive ICU transfers to identify potential predictors of deterioration. The presence of at least 1 of the following 5 risk factors was found in each case: (1) family concern about patient safety, (2) high-risk therapies including unfamiliar therapies on the unit (eg, insulin use outside of the diabetes unit), (3) elevated PEWS of ≥ 5 , (4) watcher or a patient where a clinician had a “gut feeling” that the patient was at risk for deterioration or “close to the edge,” and (5) communication concern that may impact patient safety.

Intervention

The SA intervention included the following: (1) a formalized process where bedside nurses proactively identified these 5 factors, (2) unit-based huddles where charge nurses and physicians discussed identified factors and developed mitigation plans, (3) initiation of 3-times daily inpatient huddle where individual patient risk was discussed and specific predictions made, (4) development of a continuous learning system to evaluate SA and UNSAFE transfers, and 1 year later (5) development of a “robust” and explicit plan for patients identified as having 1 of the risk factors. Figure 1 provides a model of communication and action pathway for identification of patient risk. Figure 2 is the key driver diagram that illustrated the study team's belief in hypothesized drivers needed to

achieve the aim. Table 1 provides details on individual interventions.

Proactive Identification of Risk

In our SA model (Fig 1), the fundamental job of the bedside nurses and interns was to identify any of the 5 risk factors. These clinicians have the most touch time with the patients but are generally the least experienced on the team. Tools were developed to support bedside nurses in identifying risk factors during their routine assessments. Structured yes/no questions (Fig 3) were prototyped, tested, and later spread.

Unit-Based Huddles

Unit-based huddles between the charge nurse and bedside nurse, regardless of identified risk, were scheduled every 4 hours. Huddles also were promoted whenever new risk factors were identified. The aim of the huddle was to trigger a bedside evaluation by experienced nurses and physicians of any patient with the identified risk. Huddles were led by a watchstander charge nurse and senior resident when risk was identified. The term “watchstander” was borrowed from the military to highlight that the primary job of the charge nurse

and senior resident was to know which patients were at high risk for deterioration. When concerns were identified, the clinical care team discussed the risk at the patient’s bedside and developed a plan to mitigate that risk. This provided a standardized opportunity for more experienced clinicians to coach those less experienced in both patient management and communication/escalation techniques.

Three-Times Daily Inpatient Huddle

Our design later leveraged existing structures to proactively escalate identified and unresolved safety risks. Historically, our organization conducted an 8 AM huddle that brought together a charge nurse from each inpatient unit and the manager of patient services (MPS) who oversaw the flow and staffing of inpatients. Before our work, there was no standardized discussion of patient safety. We piloted with 4 units and then spread to all noncritical care units a structured process where the charge nurse from each unit (1) reported on any risk factors present that were not fully addressed and (2) predicted any MRT activations. This process continued to be facilitated by the MPS, and a safety officer of the day (SOD) attended each inpatient huddle. The SOD was an experienced pediatrician who was given the authority by our hospital leadership to assist with mitigating any safety and/or communication concerns on the inpatient units. The MPS and SOD provided coaching on how to address concerns raised by the primary clinical team, including positive reinforcement of key behaviors and role modeling. The details of testing similar huddles at 4:30 PM and midnight are displayed in Table 1.

Continuous Learning System to Evaluate SA

The final component of our planned intervention was to develop a data




FIGURE 1

Identify, mitigate, and escalate model illustrates which risk factors were systematically identified and how standardized communication about risk occurred throughout the center.




FIGURE 2

Key driver diagram illustrates the drivers (at right) that would lead to aim through improved situation awareness and no unrecognized clinical deterioration.

TABLE 1 Specific Interventions, Settings, and Timing for Each Intervention

Category of Intervention	Specific Interventions	Setting	Timing
Proactive identification of risk	Risk categories prototyped and tested on small scale	1 general pediatric unit	November 2009
	Algorithm developed and tested throughout unit	1 general pediatric unit	December 2009
	Algorithm tested and adapted on different patient populations	4 units including subspecialty and surgical care	January 2010
Unit-based huddles	Algorithm posted and spread throughout hospital	All acute care units	March 22, 2010
	Huddle tested on small on 1 unit	1 general pediatric unit	November 2009
	Adapted to include residents only when risk identified	1 general pediatric unit	December 2009
	Tested and adapted on 4 units	4 units including subspecialty and surgical care	January 2010
	Didactic and case-based education for charge nurses	Conference room	February and March 2010
Three-times daily inpatient huddle	Spread throughout hospital	All acute care units	March 22, 2010
	Safety officer attends and each charge nurse lists any patient risks that were not fully addressed with predicted discharges and admissions	4 test units at 8 AM	January 2010
	Safety officer and MPS round on each unit	4 test units at 4 PM	January 2010
	MPS rounds on each test unit	4 test units at 12 AM	January 2010
	3-time daily inpatient huddle extends to all units	All acute care units	March 22, 2010
	Explicit predictions for calls of medical response team made	All acute care units	April 2010
	Afternoon rounds moved to huddle in conference room with each charge nurse in attendance	All acute care units	October 2010
	Overnight rounds moved to huddle in conference room	All acute care units	January 2012
	ACA form prototyped	4 units including subspecialty and surgical care	January 2010
	Weekly report with rates of risk identification and escalation and UNSAFE transfers along with patient-level story about situation awareness	All acute care units	March 2010
Continuous learning system	Control plan developed for identifying and acting on special cause with process measures and UNSAFE transfers	Used on all acute care units as needed	July 2010
	Database that combined data from ACA forms and admit/transfer data from EHR developed and tested and went into production	Used by Manager, Patient Services to track patients on all acute care units	September 2010
	Robust plan checklist generated and tested by 1 charge nurse	1 unit that specialized in transitional tracheostomy and ventilator care	March 2011
	Checklist adapted and tested with all nurses on unit	Same transitional care unit	April 2011
	Physician event note template created and tested in EHR	Same transitional care unit	June–July 2011
	Identified risk factors placed in EHR in format to be scanned by safety officer and other leaders	1 neurosciences unit	July–August 2011
	Checklist, template, and risk factors in EHR spread	All acute care units	September 2011
Robust plan			

system to rapidly identify process and outcome failures and direct this information to project leaders and leaders on individual units. To achieve this aim, (1) apparent cause analysis (ACA) forms were completed within 1 hour of each floor to ICU transfer to identify potential UNSAFE transfers and associated process failures, (2) a password-protected database was constructed to integrate information from these forms and the EHR, (3) process and outcome data were distributed each week to unit level clinical and medical directors with a story of patient-level SA, and (4)

a control plan was designed with inpatient leaders to identify special cause on tracked process and outcome measures and target further interventions.

Robust Plan

One year after SA work began, the improvement team worked with 1 inpatient unit to develop and test a checklist tool to improve the mitigation/escalation process for patients with identified risk (Table 1). During multidisciplinary discussions, we proposed a “robust plan” bundle which

included the following: (1) plan with proposed treatment change, (2) explicit communication with care team, (3) prediction of expected outcome, (4) outcome deadline, and (5) escalation plan (usually the MRT or discussion with SOD/MPS) if outcome was not achieved by a predefined deadline. This tool was tested, adapted, and spread throughout the remaining inpatient units (Fig 4). Subsequent testing integrated risk identification within the EHR and added focused discussion of a robust plan during safety rounds.




FIGURE 3 Situation awareness algorithm illustrates the tool used during education and early phases and the specific questions and communication pathways.

Study of the Intervention

In our observational time series study, data were collected on process measures of systematic identification, mitigation, and escalation of risk that we believed would improve SA and decrease UNSAFE transfers and SSEs. To evaluate the consistency of huddles and how well the identify, mitigate, and escalate intervention was implemented, data initially were collected from each unit on each nursing shift to measure the reliability that each shift identified all patients at risk and mitigated or escalated that risk. This was captured through a checklist-based form that followed the flow of algorithm of Fig 3 and was completed by each charge nurse. The tool was tested and evaluated with charge nurses from several units during early phases. Before spread throughout the hospital, 116 charge nurses received training on the process and tool through a 1.5-hour learning session. Validity of process

data were evaluated through discussions during inpatient huddles by investigators, SOD, and MPS. UNSAFE transfers were identified from the ACA process and validated against review of the EHR for each ICU transfer. SSEs were captured through a safety reporting process as previously described.²²

Analysis

Primary analysis of both process and outcome measures was performed by using statistical process control charts. For the primary outcome of UNSAFE transfers, results were tracked by using both a days-between *t*-chart and rate chart. Established rules for identifying special cause were employed.^{25–27}

RESULTS

After testing on 4 inpatient units from January 1, 2010, to March 21, 2010, the unit-level huddles and proactive

inpatient huddles began on each of the 14 noncritical care inpatient units on March 22, 2010. The process measure evaluated the consistency of huddles and specifically how frequently patient risk was identified and mitigated or escalated each nursing shift on each unit. The number of units by week where $\geq 90\%$ of weekly nursing shifts fully identified and mitigated or escalated patient risk were tracked on run charts and revealed both improved and sustained performance for 11 months of tracking (Fig 5). On each participating unit, 90% to 95% of identified risk was mitigated by the primary team with no escalation needed. Each inpatient huddle took less than 30 minutes. Although initially there was substantial variation in the number of patient risks that were escalated, a median of 2 risks for each huddle were escalated the first year. This increased over the study period with a median of 7.5 concerns escalated in May 2012.

Patient Risk Identified

↓

Risk **communicated** to Health Care Team

↓

Team develops **plan** of care with expected outcome, deadline (MAX 8 hr), and escalation plan

↓

Plan **documented** and set in motion

↓

If expected outcome is **NOT MET** in time allowed – **do escalation plan**

↓

If escalated, start a **new plan** / column

Situation Awareness Robust Planning Tool

Charge Nurse Initials _____ Unit _____ Date _____

Patient Initials			
Concern (check SA concern and describe below) 	☐ High PEWS ☐ Comm. Concern ☐ Watcher ☐ Family Concern	☐ High PEWS ☐ Comm. Concern ☐ Watcher ☐ Family Concern	☐ High PEWS ☐ Comm. Concern ☐ Watcher ☐ Family Concern
	☐ Notify MPS(Required)	☐ Notify MPS(Required)	☐ Notify MPS(Required)
Plan	Persons Notified: ☐ Respiratory Therapist ☐ Nurse Practitioner ☐ Intern ☐ Senior ☐ Fellow ☐ Attending	Persons Notified: ☐ Respiratory Therapist ☐ Nurse Practitioner ☐ Intern ☐ Senior ☐ Fellow ☐ Attending	Persons Notified: ☐ Respiratory Therapist ☐ Nurse Practitioner ☐ Intern ☐ Senior ☐ Fellow ☐ Attending
	Expected Outcome -Must be Yes / No criteria Example: HR will decrease to 120	Example: Work of breathing will decrease	Example: Family states status improved
Outcome Deadline (If more than 1 hour, state why) Example: By 8:00pm	Example: By 1:00pm	Example: By 1:00am	
Escalation Plan if outcome not met? If Escalated start new column / plan	☐ Re-consult with Senior, Fellow and/or Attending MD ☐ Call MPS ☐ Call MRT	☐ Re-consult with Senior, Fellow and/or Attending MD ☐ Call MPS ☐ Call MRT	☐ Re-consult with Senior, Fellow and/or Attending MD ☐ Call MPS ☐ Call MRT
Notes			

FIGURE 4
Situation awareness robust planning tool.

The rate of UNSAFE transfers per 10 000 non-ICU inpatient days is displayed in Fig 6. An initial decrease in UNSAFE transfers occurred, though it did not meet rules for special cause and was not sustained. Analysis of UNSAFE transfers through an ongoing ACA process revealed that in the vast majority of UNSAFE transfers, patient risk

had been identified but not fully mitigated on unit or escalated to the MRT or safety team. This led to focused improvement work on development of a robust plan as detailed above. After spread, the rate of UNSAFE transfers improved from a baseline of 4.4 to 2.4 transfers per 10 000 non-ICU inpatient days, meeting criteria for special cause

variation with 8 points below the median line (Fig 6). Additionally, a significant change in the days-between inpatient SSEs from 100 days to >400 twice was observed in association with the intervention. Shortly before this work began, the number of MRT activations and PICU transfers per month increased significantly in association

with a campaign to eliminate informal PICU consults (Fig 7). MRT-preventable codes outside the ICU were rare before and during the study period (Fig 8).

DISCUSSION

Our system to reliably identify, mitigate, and escalate patient risk was associated

with a significant decrease in UNSAFE transfer and SSEs among inpatients. By providing a proactive and reliable model to identify and rapidly intervene on newly detected risk and early patient deterioration, we improved SA across our inpatient system. SA is achieved by scanning for important information,

recognizing patterns and trends, and making short-term predictions.^{15–17} We believe that although SA is still uncommonly discussed in the medical literature, interventions such as rapid response systems and early warning scores (both in broad use) employ SA to identify deteriorating patients.²⁸ We




FIGURE 5

Process measure run chart illustrating the number of units by week where $\geq 90\%$ of weekly nursing shifts fully identified patients at risk (solid line/diamond) and where $\geq 90\%$ of weekly shifts fully mitigated or escalated that risk (dotted line/circle).




FIGURE 6

UNSAFE transfer rate chart. Rate of UNSAFE transfers per 10,000 non-ICU patient days at base location by month (n = non-ICU inpatient days by month).





FIGURE 7
Rate of MRT activations and transfers to ICU by month.



Source: Critical Care

Chart Updated through 04/30/12 by K. Simon, James M. Anderson Center for Health Systems Excellence

FIGURE 8
Rate of MRT preventable codes outside of the ICU by month.

sought to build upon this work by addressing limitations in many rapid response systems' inherently reactive (versus proactive) nature.¹⁴

We developed the concept of UNSAFE transfers to be used as the primary

outcome measure for improvement efforts with the belief that these events represented potential precursor events to serious harm such as codes and because these events occurred sufficiently often in our center to enable

rapid testing, learning, and adapting. Our time series design allowed us to learn sequentially, and we observed an initial decrease in the rate of UNSAFE transfers followed by a return to our previous baseline levels. With process

measures in place that did not reveal a decrease in reliability of our intervention, we further studied where the process failures occurred regarding UNSAFE transfers. With this data, we learned that although 1 year into the intervention we had a system that identified risk, we did not systematically address this risk. We found that even on our most high risk patients, the language of plans included terms such as “continue to observe.” Without explicit and time-bound plans, clinicians were observing patients until they received aggressive resuscitation that met criteria for UNSAFE transfer.

Our second stage of interventions testing a robust plan and prediction was designed by a multidisciplinary team of leaders and front-line physicians, nurses, and respiratory therapists. The improvement team believed that a shared mental model or team SA would not be achieved without explicit prediction and contingency planning. We believed that this was because level 3 SA (the projection of current event status in the near future) was still often not achieved. This may have been due to the limits of individual clinicians in making near-term projections (eg, this tachycardic patient will be in shock within 4 hours if we do not aggressively hydrate) but was believed to more commonly result from doctors, nurses, and other members of the care team not explicitly sharing their mental model. Our basic theory was borrowed from hypothesis testing in the scientific method and explicit predictions in plan-do-study-act cycles. The goal was to increase accountability and to make disconcerting data (eg, patient did not improve as predicted) clear to each member of the team. The spread of this intervention and its integration into proactive inpatient huddle was associated with a sustained decrease in UNSAFE transfers.

One strength of our work is that we created a system of care built on reliable

processes, not individual clinicians. We were able to build these processes into the workflows of busy clinicians and provide via the inpatient huddle a valued activity for charge nurses as they gained insight and assistance with their sickest patients. Our sustained reduction in UNSAFE transfers over the last 12 months is further evidence of the success of building interventions into work flow. Importantly, this intervention did not add additional clinicians to our system of care but instead clarified roles and processes for charge nurses, MPS, residents, and attending physicians. The additional responsibilities of SODs are on the order of 1.5 hours per day. We believe our work builds upon previous interventions to address patient deterioration such as rapid response systems and PEWS. The proactive and standardized nature of our intervention offers an important answer to afferent limb failures of the MRTs.¹⁴ Our intervention supplements the early warning score with other risk domains, most powerfully for us was that of the “watcher” or patient that a clinician has “a gut feeling is close to the edge.” We believe this employs the tacit knowledge of experienced clinicians and hence likely will achieve greater sensitivity than any numerical scoring tool, especially since we combined this concept with objective data.^{29,30} We also believe that assessment of risk as relayed by family and as emerges from communication problems has substantial face validity in identifying and predicting deterioration. Our final risk category was that of high risk therapy and borrowed from HRO thinking on the need for special oversight and procedures with new and unfamiliar therapies; for example, we believe the administration of insulin on short-stay surgery unit has a fundamentally different risk profile than doing so on diabetes unit. We therefore target these high risk

therapies and address any knowledge gaps in close to real-time. Our intervention is perhaps most similar to that of the Rover team as described by Hueckel et al.³¹ Although both are proactive in assessment of risk, meaningful differences include our intervention’s staffing model and broader scanning for risk.

Because our goal was rapid improvement of a single site, we chose a time series design, which did not allow us to address secular trends or establish causality. We believe this design was appropriate for our innovative intervention that evolved and improved through iterative quality improvement methods. This design exposed the study to potential unmeasured confounding from safety work. We do not feel this was a particularly large risk because time series data reveal the rate of inpatient SSEs had not improved in previous years’ safety work and because there were no other large interventions targeted at this population. Additionally, it is uncertain how our results would generalize to medical centers with different patient populations, staffing models, quality improvement capabilities, and safety cultures. A final limitation is that we did not have a measure of SA to reveal that this improved as an effect of our intervention. Available measures of SA involve simulated events and typically require “pausing” the event to perform assessment.³² Clearly this was not possible or ethical in course of clinical care. A recent proposed measure of SA relied on accuracy of prediction that was fundamental to our work.³³ We did use SA for much of our conceptual model, but we cannot say definitely if improved SA was the mediator between the identify, mitigate, and escalate intervention and our decreased rate of UNSAFE transfers.

Our institution additionally has applied and tested models to improve SA and

decrease adverse events in diverse clinical settings including the operating room and resident psychiatry. We also believe improved SA and systems that proactively identify, mitigate, escalate, and predict risk have applicability to other untoward events such as flow and patient/family experience failures. Next steps include tailoring and testing interventions in these settings and in other hospital systems that have different patient populations and systems of care. Although we have begun to incorporate risk factors into the EHR, we believe simulation and human factors methods are needed to evaluate the role of technology and automation in

improving SA and facilitating clinician workflows.

CONCLUSIONS

A reliable system to identify, mitigate, and escalate risk can be implemented in a children's hospital and is associated with a reduction in safety events in a context where these events were already uncommon. We believe HRO thinking and SA are potentially transformative concepts for health care systems. Models to identify risk early and reliably intervene are likely generalizable both to different clinical systems and to modify different outcomes

such as the patient/family experience and patient flow.

ACKNOWLEDGMENTS

The authors thank Ralph Brueggemann, Ron Lu, and Ripal Soni for their expertise and work in database development; Kathy Dressman for her leadership with the robust plan project; Dick Gardner, Amy Bunker, and Nancy Burse for their work in developing and facilitating education; and Pam Kiessling, Mary Jo Giaccone, Ken Tegtmeyer, and Jan Jacob for their leadership and thinking on this work. Dr Brady also thanks Tom DeWitt for his exceptional mentorship.

REFERENCES

1. Chan PS, Jain R, Nallmothu BK, Berg RA, Sasson C. Rapid response teams: a systematic review and meta-analysis. *Arch Intern Med*. 2010;170(1):18–26
2. Brill R, Gibson R, Luria JW, et al. Implementation of a medical emergency team in a large pediatric teaching hospital prevents respiratory and cardiopulmonary arrests outside the intensive care unit. *Pediatr Crit Care Med*. 2007;8(3):236–246, quiz 247
3. Sharek PJ, Parast LM, Leong K, et al. Effect of a rapid response team on hospital-wide mortality and code rates outside the ICU in a children's hospital. *JAMA*. 2007;298(19):2267–2274
4. Van Voorhis KT, Willis TS. Implementing a pediatric rapid response system to improve quality and patient safety. *Pediatr Clin North Am*. 2009;56(4):919–933
5. Hillman KM, Bristow PJ, Chey T, et al. Duration of life-threatening antecedents prior to intensive care admission. *Intensive Care Med*. 2002;28(11):1629–1634
6. Hillman K, Chen J, Cretikos M, et al; MERIT study investigators. Introduction of the medical emergency team (MET) system: a cluster-randomised controlled trial. *Lancet*. 2005;365(9477):2091–2097
7. Hanson CC, Randolph GD, Erickson JA, et al. A reduction in cardiac arrests and duration of clinical instability after implementation of a paediatric rapid response system. *Postgrad Med J*. 2010;86(1015):314–318
8. Hodgetts TJ, Kenward G, Vlackonikolis I, et al. Incidence, location and reasons for avoidable in-hospital cardiac arrest in a district general hospital. *Resuscitation*. 2002;54(2):115–123
9. Hunt EA, Zimmer KP, Rinke ML, et al. Transition from a traditional code team to a medical emergency team and categorization of cardiopulmonary arrests in a children's center. *Arch Pediatr Adolesc Med*. 2008;162(2):117–122
10. Tibballs J, Kinney S. Reduction of hospital mortality and of preventable cardiac arrest and death on introduction of a pediatric medical emergency team. *Pediatr Crit Care Med*. 2009;10(3):306–312
11. Jones DA, DeVita MA, Bellomo R. Rapid-response teams. *N Engl J Med*. 2011;365(2):139–146
12. Kotsakis A, Lobos AT, Parshuram C, et al; Ontario Pediatric Critical Care Response Team Collaborative. Implementation of a multicenter rapid response system in pediatric academic hospitals is effective. *Pediatrics*. 2011;128(1):72–78
13. Zenker P, Schlesinger A, Hauck M, et al. Implementation and impact of a rapid response team in a children's hospital. *Jt Comm J Qual Patient Saf*. 2007;33(7):418–425
14. DeVita MA, Smith GB, Adam SK, et al. "Identifying the hospitalised patient in crisis"—a consensus conference on the afferent limb of rapid response systems. *Resuscitation*. 2010;81(4):375–382
15. Endsley MR. Measurement of situation awareness in dynamic systems. *Hum Factors*. 1995;37(1):65–84
16. Endsley MR. Toward a theory of situation awareness in dynamic systems. *Hum Factors*. 1995;37(1):32–64
17. Endsley MR, Garland DJ. *Situation Awareness: Analysis and Measurement*. Mahwah, NJ: Lawrence Erlbaum Associates; 2000
18. Nadkarni VM, Larkin GL, Peberdy MA, et al; National Registry of Cardiopulmonary Resuscitation Investigators. First documented rhythm and clinical outcome from in-hospital cardiac arrest among children and adults. *JAMA*. 2006;295(1):50–57
19. Silber JH, Kaestner R, Even-Shoshan O, Wang Y, Bressler LJ. Aggressive treatment style and surgical outcomes. *Health Serv Res*. 2010;45(6 pt 2):1872–1892
20. Weick KE, Sutcliffe KM. *Managing the Unexpected: Resilient Performance in an Age of Uncertainty*, 2nd ed. San Francisco, CA: Jossey-Bass; 2007
21. US Agency for Healthcare Research and Quality, Lewin Group. *Becoming a High Reliability Organization: Operational Advice for Hospital Leaders*. Rockville, MD: Agency for Healthcare Research and Quality; 2008
22. Muething SE, Goudie A, Schoettker PJ, Donnelly LF, Goodfriend MA, Bracke TM, et al. Quality improvement initiative to reduce serious safety events and improve patient safety culture. *Pediatrics*. 2012;130(2). Available at: www.pediatrics.org/cgi/content/full/130/2/e423
23. Bonafide CP, Roberts KE, Priestley MA, et al. Development of a pragmatic measure for evaluating and optimizing rapid

- response systems. *Pediatrics*. 2012;129(4). Available at: www.pediatrics.org/cgi/content/full/129/4/e874
24. Tucker KM, Brewer TL, Baker RB, Demeritt B, Vossmeier MT. Prospective evaluation of a pediatric inpatient early warning scoring system. *J Spec Pediatr Nurs*. 2009;14(2):79–85
25. Shewhart WA, Deming WE. *Statistical method from the viewpoint of quality control*. Washington, DC: The Graduate school, the Department of agriculture; 1939.
26. Shewhart WA. *Economic Control of Quality of Manufactured Product*. New York, NY: D. Van Nostrand Company, Inc; 1931
27. Wheeler DJ. *Understanding Variation: the Key to Managing Chaos*. Knoxville, TN: SPC Press; 1993
28. VandenBerg SD, Hutchison JS, Parshuram CS; Paediatric Early Warning System Investigators. A cross-sectional survey of levels of care and response mechanisms for evolving critical illness in hospitalized children. *Pediatrics*. 2007;119(4). Available at: www.pediatrics.org/cgi/content/full/119/4/e940
29. Crandall B, Getchell-Reiter K. Critical decision method: a technique for eliciting concrete assessment indicators from the intuition of NICU nurses. *ANS Adv Nurs Sci*. 1993;16(1):42–51
30. Fackler JC, Watts C, Grome A, Miller T, Crandall B, Pronovost P. Critical care physician cognitive task analysis: an exploratory study. *Crit Care*. 2009;13(2):R33
31. Hueckel RM, Turi JL, Cheifetz IM, Mericle J, Meliones JN, Mistry KP. Beyond Rapid Response Teams: Instituting a “Rover Team” Improves the Management of At-Risk Patients, Facilitates Proactive Interventions, and Improves Outcomes. In: Henriksen K, Battles JB, Keyes MA, Grady ML, editors. *Advances in Patient Safety: New Directions and Alternative Approaches (Vol 3: Performance and Tools)*. Rockville (MD); 2008
32. Wright MC, Taekman JM, Endsley MR. Objective measures of situation awareness in a simulated medical environment. *Qual Saf Health Care*. 2004;13(suppl 1):i65–i71
33. Reader TW, Flin R, Mearns K, Cuthbertson BH. Team situation awareness and the anticipation of patient progress during ICU rounds. *BMJ Qual Saf*. 2011;20(12):1035–1042

(Continued from first page)

Dr Brady had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis; Drs Brady, Muething, and Kotagal, Mr Ashby, and Dr Wheeler contributed to the study concept and design; Drs Brady, Muething, and Kotagal, Mr Ashby, Ms Gallagher, Ms Hall, Ms Goodfriend, Dr White, Ms Bracke, Ms DeCastro, Ms Geiser, Ms Simon, Ms Tucker, Mr Olivea, and Dr Wheeler contributed to improvement project leadership and execution; Drs Brady and Ashby, Ms Gallagher, Ms Hall, Dr White, and Ms Bracke contributed to the acquisition of data; Dr Brady and Ms Bracke contributed to the statistical analysis; Drs Brady, Muething, and Kotagal, Mr Ashby, Ms Gallagher, Ms Hall, and Drs Conway and Wheeler contributed to the interpretation of data. Drs Brady and Wheeler drafted the article; and all authors contributed to critical revision of the article for important intellectual content.

www.pediatrics.org/cgi/doi/10.1542/peds.2012-1364

doi:10.1542/peds.2012-1364

Accepted for publication Jul 26, 2012

Address correspondence to Patrick Brady, MD, MSc, Cincinnati Children's Hospital Medical Center, ML 9016, 3333 Burnet Ave, Cincinnati, OH 45229. E-mail: patrick.brady@cchmc.org

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2013 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they have no financial relationships relevant to this article to disclose.

FUNDING: Dr Brady was supported by funds from the Bureau of Health Professions, Health Resources and Services Administration, Department of Health and Human Services, under grant T32 HP10027. Portions of this project were supported by cooperative agreement number U19 HS021114 from the Agency for Healthcare Research and Quality.

Improving Situation Awareness to Reduce Unrecognized Clinical Deterioration and Serious Safety Events

Patrick W. Brady, Stephen Muething, Uma Kotagal, Marshall Ashby, Regan Gallagher, Dawn Hall, Marty Goodfriend, Christine White, Tracey M. Bracke, Victoria DeCastro, Maria Geiser, Jodi Simon, Karen M. Tucker, Jason Olivea, Patrick H. Conway and Derek S. Wheeler

Pediatrics 2013;131:e298; originally published online December 10, 2012;

DOI: 10.1542/peds.2012-1364

Updated Information & Services	including high resolution figures, can be found at: http://pediatrics.aappublications.org/content/131/1/e298.full.html
References	This article cites 25 articles, 8 of which can be accessed free at: http://pediatrics.aappublications.org/content/131/1/e298.full.html#ref-list-1
Citations	This article has been cited by 4 HighWire-hosted articles: http://pediatrics.aappublications.org/content/131/1/e298.full.html#related-urls
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Administration/Practice Management http://pediatrics.aappublications.org/cgi/collection/administration:practice_management_sub Quality Improvement http://pediatrics.aappublications.org/cgi/collection/quality_improvement_sub
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://pediatrics.aappublications.org/site/misc/Permissions.xhtml
Reprints	Information about ordering reprints can be found online: http://pediatrics.aappublications.org/site/misc/reprints.xhtml

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2013 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

