

SYSTEMATIC REVIEW

Open Access



A systematic review of adaptations of evidence-based public health interventions globally

Cam Escoffery^{1*} , E. Lebow-Skelley¹, R. Haardoerfer¹, E. Boing¹, H. Udelson¹, R. Wood², M. Hartman², M. E. Fernandez² and P. D. Mullen²

Abstract

Background: Adaptations of evidence-based interventions (EBIs) often occur. However, little is known about the reasons for adaptation, the adaptation process, and outcomes of adapted EBIs. To address this gap, we conducted a systematic review to answer the following questions: (1) What are the reasons for and common types of adaptations being made to EBIs in community settings as reported in the published literature? (2) What steps are described in making adaptations to EBIs? and (3) What outcomes are assessed in evaluations of adapted EBIs?

Methods: We conducted a systematic review of English language publications that described adaptations of public health EBIs. We searched Ovid PubMed, PsycINFO, PsycNET, and CINAHL and citations of included studies for adapted public health EBIs. We abstracted characteristics of the original and adapted populations and settings, reasons for adaptation, types of modifications, use of an adaptation framework, adaptation steps, and evaluation outcomes.

Results: Forty-two distinct EBIs were found focusing on HIV/AIDS, mental health, substance abuse, and chronic illnesses. More than half (62%) reported on adaptations in the USA. Frequent reasons for adaptation included the need for cultural appropriateness (64.3%), focusing on a new target population (59.5%), and implementing in a new setting (57.1%). Common adaptations were content (100%), context (95.2%), cultural modifications (73.8%), and delivery (61.9%). Most study authors conducted a community assessment, prepared new materials, implemented the adapted intervention, evaluated or planned to evaluate the intervention, determined needed changes, trained staff members, and consulted experts/stakeholders. Most studies that reported an evaluation ($k = 36$) included behavioral outcomes (71.4%), acceptability (66.7%), fidelity (52.4%), and feasibility (52.4%). Fewer measured adoption (47.6%) and changes in practice (21.4%).

Conclusions: These findings advance our understanding of the patterns and effects of modifications of EBIs that are reported in published studies and suggest areas of further research to understand and guide the adaptation process. Furthermore, findings can inform better reporting of adapted EBIs and inform capacity building efforts to assist health professionals in adapting EBIs.

Keywords: Adaptation, Intervention, Modifications, Implementation, Evidence-based

* Correspondence: cescoff@emory.edu

¹Rollins School of Public Health, Emory University, 1518 Clifton Road, Atlanta, GA 30322, USA

Full list of author information is available at the end of the article



© The Author(s). 2018 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Background

Emphasis on evidence-based practice in medicine, public health, and the social services has led to a prominence of the application of practice guidelines and evidence-based interventions or EBIs. When situating an EBI in a new context, public health professionals, or health practitioners who work in community settings, sometimes adapt the EBI during the process of replication [1, 2]. However, in planning and implementing these interventions, there may be mismatches between the original EBI and the characteristics of the population of interest, implementing agency, and/or community [3]. In addition, agencies may lack the resources, funding, or expertise to deliver the EBI as it was originally intended [4]. Consequently, public health professionals often make both intended and unplanned program adaptations to the EBI to better fit the new audience or context.

The concept of program adaptation was originally introduced by Rogers when he defined adaptation as the degree to which an innovation is modified in the process of its adoption and implementation [5]. Other definitions have evolved in the era of translation of EBIs and the emergence of adaptation frameworks (Table 1). These definitions share

similar characteristics, including modifying a program to meet the needs of the target population, local circumstances, or new contexts. Some definitions focus on the need to retain the core components or logic of the program [6–11]. The adaptations could be deletions, additions, or modifications [11]. Some posit that adaptations should be systematic or planned [12–14] to involve stakeholder input and to have a more rigorous process in program planning, while the CSAP (Center for Substance Abuse Prevention) framework notes that adaptations could be accidental modifications [11]. Moore and colleagues proposed the schema of timing of adaptation of proactive (planned) vs. reactive [15]. Furthermore, three definitions included modifications related to matching the culture for the new population, “cultural adaptation” [11, 16, 17]. A few definitions specify elements that could be changed such as program components, content, provider, and delivery [11, 14]. Of these definitions, CSAP’s Guidelines for Adaptation [11], Map of Adaptation Process [16, 18], ADAPT-ITT [9], and Research-based Program Adaptation [6] are cited most frequently in the published literature. In summary, although many adaptation definitions share similar characteristics, the most frequently cited ones do

Table 1 Definitions of adaptation

Article	Adaptation definition
Backer (CSAP, 2002) [11]	“The deliberate or accidental modification of the program, including the following: a. Deletions or additions (enhancements) of program components; b. Modifications in the nature of the components that are included; c. Changes in the manner or intensity of administration of program components called for in the program manual, curriculum, or core components analysis; or d. Cultural and other modifications required by local circumstances.”
McKleroy et al. 2006 [16]	Quotes Rogers’ (1995) definition and the CSAP definition (see above).
Solomon et al. [6]	Modifying an efficacious program to meet the needs of its new target population and community context while retaining fidelity (or adherence) to its core components.
Smith and Caldwell [14]	“Evidence-based programs should not be changed randomly but should be modified based on a careful review of program content, the theoretical underpinnings involved, and the context of the new environment. Four different forms of adaptation need to be considered: structural, content, provider, and delivery.”
Wingood and DiClemente [9]	“The process of modifying an EBI without competing with or contradicting its core elements or internal logic.”
Barrera and Castro, Kumpfer et al. [17, 22]	Developing cultural adaptations or accommodations of EB practices for international transport is a ... “process requiring careful assessment of the local political, religious, and economic context as well as the cultural norms and family practices of country and internal ethnic groups. It should be a careful and rigorous process ...guided by research and theory.”
Lee et al. [7]	“Inherent in [the process of moving evidence-based programs (EBPs) from research to practice] is the tension between implementing programs with fidelity and the need to tailor programs to fit the target population.”
Card et al. [3]	“The process of altering a program to reduce mismatches between its characteristics and those of the new context in which it is to be implemented or used.”
Chen et al. [8]	“Methods of planned adaptation identify differences in the new target population and attempt to make changes to the EBI that accommodate these differences without diluting the program’s effectiveness.”
Rolleri [10]	“The process of making changes to a program in order to make it more suitable for a particular population or for an organization, based upon its capacity. Changes to a program should be made without compromising or deleting the program’s core components.”
Bartholomew et al. [42]	Systematic adaptation requires that planners make adaptation decisions by comparing the logic of change in the EBI with the needs of the new community. Planners should only make changes that correspond with mismatches between the EBI and community needs.

not emphasize the same concepts. Thus, it is important to discern how professionals in the field describe their adaptations, why they make modifications and the types of changes that they make, and how they use frameworks to conduct adaptations.

Previous reviews have found that modifications to original EBIs often occur spontaneously when they are adopted into other practice settings [1, 15]. Common reasons for adaptations include responding to participants' attributes [18, 19], needs [20] or culture [15], constraints such as limited time or resources [15, 19–21], issues related to participant recruitment or retention [15], and accommodating practice or setting circumstances/context [20].

Increased development of models and frameworks to guide the adaptation of EBIs began with national EBI dissemination efforts related to disease prevention areas in substance abuse and HIV/AIDS [6, 9, 14] or cultural adaptations to existing programs [7, 22]; these frameworks provide approaches to facilitate adaptation. Escoffery and associates recently conducted a scoping study that found 13 adaptation frameworks [23]. They reported 11 common steps including assess the community, understand the intervention, select intervention, consult with experts, consult with stakeholders, decide what needs adaptation, adapt the original program, train staff, test the adapted materials, implement, and evaluate. These frameworks enhance the translation of evidence-based practices. As Wandersman's Interactive Systems Framework suggests, supports are necessary to guide the public health system or agencies to adopt and implement new public health interventions [24]. These frameworks assist public health professionals as capacity building tools for a structured adaptation process.

Limited research has explored *how* adaptation occurs in practice. Little is understood about who is involved in adaptation processes, what common types of changes are made to the original program, and what mechanisms are used. This review advances the concept of adaptation and elucidates common adaptation processes in real-world community settings as reported in the published literature. Community settings are defined as various organizations or places in communities such as schools, faith-based organizations, social services or public health agencies, households, or worksites. The research questions for the review were as follows:

1. What are the reasons for and common types of adaptations being made public health EBIs in community settings as reported in the published literature?
2. What steps are described to making adaptations to EBIs?
3. What outcomes are assessed in evaluations of adapted EBIs?

Methods

We followed procedures for systematic reviews based on the Cochrane Handbook of Systematic Reviews of Public Health Interventions [25] and the reporting guideline, Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [26].

Search strategy

We searched Ovid PubMed, PsycINFO, PsycNET, and CINAHL with the assistance of an experienced health sciences librarian. The date of the last search was December 2015. Concepts for the search included adaptation, evidence-based interventions and practice, health behavior, and quality of healthcare. Combinations of the associated MeSH terms were used to develop the initial PubMed search and then adapted to search other databases. The search strategies can be found in Additional file 1. We also manually cross-referenced reference lists of included studies. We downloaded relevant citations into a reference manager software program, EndNote, which facilitated removing duplicate citations identified in the multiple databases. We exported the resulting composite library into an Excel file for documentation of the title and abstract review process.

Eligibility criteria

The project team created an Eligibility Assessment Checklist restricting included articles to those reporting primary studies published in English after 1995 and that examined the adaptation process or outcomes of an adapted evidence-based intervention (public health program or policy). Programs are defined as a combination of strategies designed to create behavior change or improve health status and policies are rules, regulations, or actions related to a health goal or service. These adaptations reported could be proactive (purposeful) or reactive. Articles were excluded if they did not describe the adaptation methods or if the full-text article was unable to be located after an exhaustive search. We combined articles reporting different aspects of the same EBI, e.g., the evaluation findings and the adaptation process.

Screening

Two trained reviewers (CE, HU) independently screened titles and abstracts after duplicates were removed, using the Eligibility Assessment Checklist. We selected potentially relevant abstracts for a full-text review conducted independently by the two trained reviewers. The first author resolved any disagreement between the reviewers.

Study quality assessment

We assessed study quality of the articles based on their use of a theory or adaptation framework, and in the case of those that included an evaluation, we assessed the

rigor of the design. We used these variables descriptively, however, and did not differentiate studies based on these variables.

Data abstraction and analysis

We reviewed the articles of EBI adaptations for six categories of variables: (1) characteristics of the original and adapted EBI (name, disease/topic, population and setting), (2) reason(s) for adaptation, (3) type(s) of modifications, (4) steps (tasks) in adaptation described by the authors, (5) reference to an adaptation framework, and (6) measures of implementation and intervention outcomes (see definitions in Additional file 2). In addition, we described how they presented the EBI adaptations made in the articles. We used a structured data abstraction form designed in Excel 2016 to record the extracted information. We used Stirman and associates' typology of modifications [1], the adaptation steps identified in the scoping study of adaptation frameworks [23], and implementation outcomes defined by Proctor and colleagues [27]. Context modifications were defined as changes to format, location, or personnel delivering the intervention, while content modifications were changes to the intervention materials, procedures, or delivery. In coding adaptation steps, we combined consulting with stakeholders and experts and had an "other" option, resulting in nine named steps. For each study, we examined bias in the study through documentation of participants (e.g., selection, generalizability), study design, and inadequate results reporting. Two trained reviewers (CE, HE, RW, ME, PDM, EL) independently coded the included articles. Discrepancies were discussed and adjudicated by the larger team.

Data synthesis and presentation


We presented summaries of study-specific adaptation reasons, steps, types of adaptations, and outcome measures with descriptive statistics across studies. We described the original and adapted EBI, the study population, reasons for adaptations, the name of adaptation frameworks, and examples of adaptations qualitatively.

Results

We found 543 unique citations that yielded 45 articles reporting 42 distinct program adaptations after the two levels of screening (Fig. 1). Main reasons for exclusion were a lack of description of the adaptation process or methods, not being a public health program or policy, and not being a primary study (e.g., protocol, review).

Adaptation characteristics

Publication years of the primary citations are from 2003 to 2014, and common disease topics included HIV/AIDS, mental health, substance abuse, and chronic



illnesses (Table 2). Many of these studies had non-experimental designs ($k = 27$, 64.3%), and the remainder had experimental ($k = 12$, 28.6%) or quasi-experimental ($k = 3$, 7.1%) designs. Thirty-six adaptations included an evaluation. Most ($k = 26$, 61.9%) reported on adaptations that took place in the USA, and one EBI was adapted in three locations (USA, Africa, and Asia). Other EBIs were adapted in Africa ($k = 4$), Asia ($k = 5$), Europe ($k = 3$), Canada ($k = 1$), the Caribbean ($k = 1$), and Australia ($k = 1$).

Reasons for adaptation

The most common reasons for adaptation included the need for a culturally appropriate program ($k = 27$; 64.3%), a new target population ($k = 25$; 59.5%), and a new community setting ($k = 24$; 57.1%) (Table 2). Less common reasons for adaptation were the desire to improve ease and feasibility of implementation ($k = 6$; 14.3%), attempting to make the program more widely accessible ($k = 1$; 2.4%), and trying to condense the original intervention ($k = 1$; 2.4%).

Types of modifications

Authors reported making an average of 3.4 (SD = 0.90, range 2–5) different types of adaptations with a mode of 3 (Table 3). All 42 (100%) reported some modification of the EBI content. The form this took usually included tailoring ($k = 39$; 92.9%) or adding elements ($k = 30$; 71.4%). For example, Cornelius and associates modified HIV prevention videos originally tested with young pregnant women to be relevant to older African American

Table 2 Characteristics of included reports, reasons for adaptation, and frameworks

First author, year	Original EBI name (adapted EBI name) study design	EBI disease/topic	Reason for adaptation	Target population/setting		Adaptation framework (if mentioned)
				Original (author)	Adapted	
Reijneveld, 2003 [53]	Healthy & Vital (no change) Experimental—RCT	Physical inactivity/poor physical and mental well-being	Cultural appropriateness New population	People aged ≥ 65 in welfare services in the Netherlands	Turkish immigrants aged ≥ 45 in welfare services in the Netherlands	N/R
Komro, 2004 [54]	Project Northland (Project Northland Chicago) Experimental—RCT	Alcohol use	Cultural appropriateness New population New community	Mostly white, 6th–8th grade students in rural NE Minnesota, USA	Culturally diverse 6th–8th grade students in Chicago, IL, USA	N/R
Sarkisian, 2005 [55]	Empowerment: Facilitating a Path to Personal Self-Care (N/R) Non-experimental	Diabetes	Cultural appropriateness New population	Younger, mostly Caucasian patients with diabetes in the USA	African Americans and Latinos aged ≥ 55 with diabetes in public health diabetes and geriatrics clinics and senior centers in South Los Angeles, CA, USA	N/R
Tsey, 2005 [56]	Family Wellbeing (no change) Non-experimental	Teasing, bullying, fighting, low self-esteem, truancy	New population New community	Adults in Aboriginal Australia (Tsey [57])	Students in 2 primary schools in remote indigenous communities in Cape York Peninsula, far north Queensland, Australia	N/R
Villarruel, 2005 [58]	Be Proud! Be Responsible! (iCuideat) Experimental—RCT	HIV	Cultural appropriateness New population	African American adolescents aged 13–18 from community-based agencies in Philadelphia, USA	Inner-city Latino adolescents aged 10–19 in Philadelphia, PA, USA	N/R
Belanksy, 2006 [59]	Integrated Nutrition Education Program, INP (Integrated Nutrition and Physical Activity Program, INPAP) Non-experimental	Nutrition and physical activity	Cultural appropriateness New community	Elementary school children in a school setting in Denver, USA	2nd and 3rd grade students in a rural, biethnic, low-income county in south-central Colorado, USA	N/R
Hitt, 2006 [60]	Project RESPECT (N/R) Non-experimental	HIV/STD	New population New community	Heterosexual individuals aged ≥ 14 attending 5 public STD clinics in the USA (Kamb [61])	MSM, IDU and heterosexual individuals attending either a local health department or a CBO for prevention counseling services in Texas, USA	N/R
Somerville, 2006 [62]	Popular Opinion Leader, POL (Young Latino Promotors, YLP) Non-experimental	HIV	Cultural appropriateness New population	White gay men in gay venues frequented predominantly by whites in mid-sized southern cities in the USA	Latino migrant MSM aged 18–30 in Texas and California USA-Mexico border communities	N/R

Table 2 Characteristics of included reports, reasons for adaptation, and frameworks (Continued)

First author, year	Original EBI name (adapted EBI name) study design	EBI disease/topic	Reason for adaptation	Target population/setting		Adaptation framework (if mentioned)
				Original (author)	Adapted	
NIMH Collaborative HIV/STD Prevention Trial Group, 2007 [38]	Community Popular Opinion Leader (C-POL) (no change) Experimental—RCT	HIV/STD	Cultural appropriateness New community	Populations vulnerable to HIV risk behavior in the USA	Individuals aged 18–49 at food markets with individually owned stalls in Fuzhou, China and individuals aged 18–30 in the following settings: wine shops in slums in Chennai, India; gathering points of young, high-risk people in barrios in Lima, Chiclayo, and Trujillo, Peru; trade school dorms in St. Petersburg, Russia; and retail establishments in rural Zimbabwe	N/R
Tsarouk, 2007 [63]	Reconnecting Youth (RY) (no change) Non-experimental	Substance abuse and HIV transmission	Cultural appropriateness	High-risk students aged 14–18 in the USA	Russian adolescents aged 14–17 with poor school performance and mild behavioral problems in schools in Moscow, Russia	N/R
Beattie, 2008 [64]	Swim and Survive, and Infant Aquatics (Water Safety in the Bush) Non-experimental	Water safety/drowning	Cultural appropriateness New community	Infants, children 5–14 years, and parents in Australia	Children and adults in rural and remote Australian communities	N/R
Cornellus, 2008 [28]	Sisters Informing Sisters on Topics about AIDS, SISTA (Women Informing Women on Topics about AIDS, WWITA) Non-experimental	HIV/AIDS	New population New community	Young African American girls in heterosexual relationships in San Francisco, USA (D'Clemente [65])	African American women ≥ 50 in heterosexual relationships who frequent churches located in low-income areas of North Carolina, USA	N/R
Gittlin, 2008 [30]	Chronic Disease Self-Management Program, CDSMP (Harvest Health, HI) Vietnamese Focus on Kids (Exploring the World of Adolescents, EWA) Non-experimental	Chronic disease self-management HIV/AIDS	New population New community Cultural appropriateness New community	Middle-class white patients aged ≥ 40 in community-based sites in the USA (Lorig [66]) Adolescents in Khanh Hoa Province, Vietnam	African Americans aged ≥ 60 with chronic condition(s) in a senior setting in Philadelphia, PA, USA Adolescents aged 15–21 in both rural and urban Vietnam	N/R N/R
Steiker, 2008 [29]	Keepin' It REAL (Refuse, Explain, Avoid, and Leave) (N/R) Quasi-experimental	Substance abuse prevention	Cultural appropriateness New population New community	Middle school youth in the USA	Adolescents aged 14–19 in high risk, unique community settings in Texas, USA	Castro - cultural adaptation
Burgio, 2009 [68]	Resources for Enhancing Alzheimer's Caregiver Health, REACH II (REACH OUT: Offering Useful Treatments) Non-experimental	Alzheimer's disease	Implementation ease/feasibility	In-home Alzheimer's caregivers in USA cities	Alzheimer's caregivers in Area Agencies on Aging in Alabama, USA	N/R
Fiscian, 2009 [69]	Making Proud Choices (N/R) Non-experimental	HIV/AIDS	Cultural appropriateness New population New community	Minority adolescents in the USA	Adolescent girls aged 10–14 in a church-affiliated junior secondary school in Ghana	N/R

Table 2 Characteristics of included reports, reasons for adaptation, and frameworks (Continued)

First author, year	Original EBI name (adapted EBI name) study design	EBI disease/topic	Reason for adaptation	Target population/setting		Adaptation framework (if mentioned)
				Original (author)	Adapted	
Mueller, 2009 [31]	<i>Cuidate!</i> (no change) Non-experimental	HIV/AIDS	New community	Latino youth aged 13–18 in Northeast Philadelphia schools, USA	Latino youth in a urban high school in Denver, CO, USA	N/R
Pekmez, 2009 [70]	Individually tailored physical activity print intervention (<i>Seamos Activas</i>) Experimental—RCT	Physical inactivity and related chronic illnesses	Cultural appropriateness New population	Sedentary adults in the USA	Overweight/obese Latinas aged 18–65 with low income and acculturation in Providence, RI, USA	N/R
Stevens, 2009 [71]	REACH II (Support Teams for Caregivers) Non-experimental	Alzheimer's disease or dementia	Implementation ease/feasibility	Family caregivers of patients with Alzheimer's disease or dementia in 5 USA cities	Family caregivers of patients with Alzheimer's disease or dementia in Texas, USA	RE-AIM and REP
DePue, 2010 [72]	Project Sugar 2, PS2 (Diabetes Care in American Samoa) Experimental—RCT	Type 2 diabetes	Cultural appropriateness New population	Urban African Americans aged ≥ 25 with diabetes in Baltimore, USA	Individuals aged ≥ 21 with type 2 diabetes in American Samoa	Lau's framework for cultural adaptation
Domenech Rodriguez, 2011 [73]	Parent Management Training—Oregon Model, PMTO (<i>Creando con Amor: Promoviendo Armonía y Superación, CAPAS</i>) Experimental—RCT	Parenting	Cultural appropriateness New population	Divorcing mothers with sons in 1st–3rd grades in a medium-sized city in the Pacific NW, USA (Forgatch [74])	Spanish-speaking Latino parents or relatives who co-parent in rural Utah, USA	CAP and EVM
Poulsen, 2010; [39] Vandenhouh, 2010 [75]	Parents Matter! (Families Matter!) Non-experimental	HIV	Cultural appropriateness New community	African American parents of preteens aged 9–12 in a controlled clinical setting in the USA	Families with children aged 9–12 in Asembo, rural west Kenya	MAP
Sadler, 2010 [76]	"Cancer Clinical Trials: The Basics" and " <i>Conversamos un rato: Información para combatir el cáncer en su comunidad</i> " (N/R) Non-experimental	Breast cancer	Cultural appropriateness New population	Individuals with cancer in the USA	African American/Hispanic American women, or women from diverse communities with breast cancer in California, USA	N/R
Rotheram-Borus, 2011 [77]	Project TALC (LA Project TALC in Los Angeles, Family to Family in Thailand, Mentor Mothers in South Africa) Non-experimental	HIV	Cultural appropriateness New community	Parents living with HIV and their children or caregiver supports in New York City, USA	Parents living with HIV and their children or caregiver supports in the USA (Los Angeles, CA), Thailand, and South Africa	CQI
Cardona, 2009 [78]	Parent Management Training—Oregon Model, PMTO (N/R) Non-experimental	Parenting/mental health	Cultural appropriateness New population New community	Divorcing mothers with sons in 1st–3rd grades in a medium-sized city in the Pacific Northwest, USA (Forgatch [74])	Latino immigrant parents with children aged 6–12 with mild behavioral problems in Detroit, MI, USA	EVM
Feinberg, 2012 [79]	Problem-Solving Treatment (Problem-Solving Education) Experimental—RCT	Depression	New population	Adults with depression in general practices in Oxford, United Kingdom (Gath [80])	Mothers with limited incomes and high rates of depression in 3 settings where they receive services in Massachusetts, USA	Backer's 6-step approach Castro's cultural adaptation
Parker, 2012; [81]	Arthritis Self-Help Program, ASHP (no change)	Arthritis	Cultural appropriateness New population Condense program	Younger, mostly non-Hispanic white adults in the USA	African American, Hispanic, and non-Hispanic white older adults attending senior centers in New York City, NY, USA	M-PACE
Chen, 2013 [82]	Non-experimental					

Table 2 Characteristics of included reports, reasons for adaptation, and frameworks (Continued)

First author, year	Original EBI name (adapted EBI name) study design	EBI disease/topic	Reason for adaptation	Target population/setting		Adaptation framework (if mentioned)
				Original (author)	Adapted	
Reid, 2012 [83]	Cognitive Behavioral Stress Management (CBSM) (no change) Non-experimental	Substance abuse, sexual behavior, and HIV	Cultural appropriateness New community	Drug abusers	HIV-positive substance abusers in recovery in Trinidad and Tobago	N/R
Rosati, 2012 [84]	Family Matters (Thai Family Matters) Experimental—RCT	Alcohol, tobacco, and other drug use	Cultural appropriateness New community	Parents and children in the USA	Adolescents aged 13–14 and their parents in Bangkok, Thailand	N/R
Tomioka, 2012 [85]	Chronic Disease Self-Management Program, CDSMP (Ke Ola Pono) Non-experimental	Chronic disease self-management	Cultural appropriateness New population	Adults aged ≥ 40 with chronic diseases in community-based sites in California, USA (Lorig [60])	Asians and Pacific Islanders with chronic diseases in Hawaii, USA	CDC's adaptation traffic light
Danielson, 2013 [83]	Sistas Informing, Healing, Living, and Empowering, SHLE (SHLEWeb) Non-experimental	HIV/STD	Cultural appropriateness Implementation ease/feasibility	African American adolescents in community health agencies in the USA (DiClemente [65])	Community-dwelling traditionally underserved African American girls aged 13–18 in the Southeast USA	N/R
Fasula, 2013 [86]	Project Safe (Project POWER) Non-experimental	HIV/STD	New population New community	African American and Mexican American women in STD clinics in San Antonio, USA (Shain [87])	HIV-negative women with sentences up to 14 months due to be released within 6 months in North Carolina women's prison facilities, USA	MAP
Parker, 2013a; [88]	Healthy Living Project (Supporting Youth and Motivating Positive Action, SYMPA) Non-experimental	HIV/AIDS	New population New community	Adults living with HIV in the USA	Youth aged 15–24 living with HIV/AIDS in Kinshasa, Democratic Republic of the Congo	ADAPT
Parker, 2013b [89]	Reciprocal imitation training (RIT) (no change) Non-experimental	ASD	Implement in new community setting Make program more widely accessible	Individuals working with children with ASD, including parents, in the USA	Individuals working with children with ASD, including parents, in the participants' homes and research lab in the Midwestern USA	N/R
Williams, 2013 [91]	Adherence Through Home Education and Nursing Assessment, ATHENA (N/R) Experimental—RCT	HIV/AIDS	Cultural appropriateness New population New community	European, African and Hispanic individuals with a high prevalence of substance abuse and mental illness for whom ARV therapy was prescribed in the northeastern USA	Patients living with HIV/AIDS receiving ARV therapy from the Hunan China CARES clinical program in rural south central China	Castro's cultural adaptation
Baydala, 2014 [92]	Life Skills Training, LST (Nimi Icinohabi) Quasi-experimental design	Substance abuse	Cultural appropriateness New population	Elementary, middle, and high school students, including ethnic minority youth in the USA	Aboriginal school-age children in Central Alberta, Canada	N/R
Broning, 2014 [93]	Strengthening Families Program for Parents and Youth 10–14, SFP 10–14 (Familien Stärken) Experimental—RCT	Substance abuse	Cultural appropriateness New community	Adolescents aged 10–14 and their caregivers in rural economically deprived regions in Iowa, USA	Adolescents aged 10–14 and their caregivers in socially deprived urban districts in Hamburg, Schwerin, Hanover and Munich, Germany	N/R
Carliou, 2014 [94]	Pool Cool (no change) Non-experimental	Skin cancer	New population Implementation ease/feasibility	Aquatics instructors, kids aged 5–10, parents and other pool users in Hawaii and Massachusetts, USA (Glanz [95])	Children and adolescents aged 2–17 enrolled in swim lessons at the Payette Municipal Pool, rural Idaho, USA	N/R

Table 2 Characteristics of included reports, reasons for adaptation, and frameworks (*Continued*)

First author, year	Original EBI name (adapted EBI name) study design	EBI disease/topic	Reason for adaptation	Target population/setting		Adaptation framework (if mentioned)
				Original (author)	Adapted	
Reback, 2014 [96]	Gay-specific cognitive behavioral therapy, GCBT (Getting Off: A Behavioral Treatment Intervention for Gay and Bisexual Male Methamphetamine Users) Experimental - RCT	Methamphetamine use/HIV	New community	Methamphetamine-using gay and bisexual men in a controlled clinical setting in the USA	Methamphetamine-using MSM in a community-based HIV prevention setting in Los Angeles, CA, USA	N/R
Riggs, 2014 [32]	Family Overweight: Comparing Use of Strategies, FOCUS (Family Wellness Program, FWP) Non-experimental	Pediatric obesity	Implementation ease/feasibility New community	Obese children and their parents in the USA (Saelens [97])	Obese children aged 6–12 and their parents in primary care clinics near Seattle, WA, USA	N/R
Tu, 2014 [98]	Clinic-based educational program to promote CRC screening among Chinese immigrants (N/R) Quasi-experimental	Colorectal cancer screening	New population Implementation ease/feasibility	Chinese immigrant in a community health center in the metropolitan area of Seattle, USA (Tu [99])	Vietnamese patients of community health centers in the metropolitan area of Seattle, WA, USA	Diffusion of innovations theory

N/R not reported, ASD autism spectrum disorder, ARV antiretroviral, CAP cultural adaptation process, CBO community-based organization, COI continuous quality improvement, EVM ecological validity model, IDU injection drug user, M-PAACE Method for Planned Adaptation through Community Engagement, MAP Map of Adaptation Process, MSM men who have sex with men, RE-AIM Reach, Effectiveness, Adoption, Implementation and Maintenance, REP Replicating Effective Programs, STD sexually transmitted disease

Table 3 Summary of adaptation characteristics reported in peer-reviewed literature (EBIs), $k = 42$

Adaptation characteristics	Studies reporting characteristic k (%)
Type of modification	
Content	42 (100%)
Tailoring	39 (92.9%)
Adding elements	30 (71.4%)
Shortening	13 (31.0%)
Removing elements	12 (28.6%)
Loosening structure	10 (23.8%)
Lengthening	9 (21.4%)
Substitution	7 (16.7%)
Integrating other approach	5 (11.9%)
Reorder elements	4 (9.5%)
Integrating intervention	2 (4.8%)
Departing	2 (4.8%)
Repeating elements	1 (2.4%)
Cultural modification	31 (73.8%)
Context	40 (95.2%)
Population	33 (78.6%)
Setting	29 (69.0%)
Other	3 (7.1%)
Delivery	26 (61.9%)
Deliverer	16 (38.1%)
Mode/medium	14 (33.3%)
Other	4 (9.5%)
Training	16 (38.1%)
Evaluation	19 (45.2%)
Change to core elements	4 (9.5%)
Reasons for adaptation	
Cultural appropriateness	27 (64.3%)
Focus on new target population	25 (59.5%)
Implement in new community setting	24 (57.1%)
Improve ease and feasibility of implementation	6 (14.3%)
Make program more widely accessible	1 (2.4%)
Condense program	1 (2.4%)
Outcomes	
Implementation	
Acceptability	28 (66.7%)
Fidelity	22 (52.4%)
Feasibility	22 (52.4%)
Adoption	20 (47.6%)
Sustainability	11 (26.2%)
Other	5 (11.9%)

Table 3 Summary of adaptation characteristics reported in peer-reviewed literature (EBIs), $k = 42$ (Continued)

Adaptation characteristics	Studies reporting characteristic k (%)
Behavioral/program	
Behavior	30 (71.4%)
Practice	9 (21.4%)
Knowledge	7 (16.7%)
Self-efficacy	5 (11.9%)
Environment	4 (9.5%)
Well-being/health	3 (7.1%)
Attitudes	3 (7.1%)
Skills	3 (7.1%)
Communication	2 (4.8%)
Policy	0
Other	4 (9.5%)
Individual satisfaction	11 (26.2%)

women [28]. In the adaptation reported by Steiker, the study team added four new videos to accompany the curriculum and rewrote scenarios used in the workbooks to incorporate local culture [29]. In the EBI adapted by Gitlin and associates, a moment of silence was added at the beginning of each session to recognize spiritual practices and their importance to participants [content modification-adding elements [30]]. More than half of the authors reported shortening the original EBI as one of the adaptations made. For the 42 programs, some teams described adapting the intervention content by shortening it ($k = 13$; 31.0%), removing elements ($k = 12$; 28.6%), loosening the structure ($k = 10$; 23.8%), lengthening the program ($k = 9$; 21.4%), substituting modules or activities ($k = 7$; 16.7%), or integrating other approaches to the intervention ($k = 5$; 11.9%).

Nearly all of the adaptations ($k = 40$; 95.2%) described modifying context, and 31 (73.8%) included cultural modifications. Most context modifications included making changes to the original EBI by adapting it to fit with the new intervention population ($k = 33$; 78.6%) and setting ($k = 29$; 69.0%). Mueller and colleagues, for example, adapted a curriculum originally delivered in community agencies and after-school programs to a school setting [31]. Over half of the adaptations included changes to the delivery of the original intervention ($k = 26$; 61.9%), either by modifying the role of the personnel delivering the intervention ($k = 16$; 38.1%) or by adapting the format or channel of delivery ($k = 4$; 33.3%). Masters-level research interventionists, for instance, delivered the family-based behavioral pediatric obesity treatment rather than medical staff in the intervention reported by Riggs and colleagues [32]. In the

EBI adapted by Danielson and team, a web-based delivery platform was used instead of small group sessions [33]. Fewer authors reported modifying procedures for training personnel ($k = 16$; 38.1%) or for evaluating the program ($k = 19$; 45.2%). Four (9.5%) studies described changing what they regarded as core elements of the original EBI.

Patterns of adaptation types

The most common combinations were content, context, and delivery ($k = 9$), and content and context ($k = 8$) (Fig. 2). Content and context were part of all other combinations; four other combinations only had one study each.

For content adaptations, the classifications reported varied greatly. However, some patterns emerged with certain content combinations, including tailoring, adding elements, and cultural modifications ($k = 5$); tailoring, adding elements, loosening structure, and cultural modifications ($k = 4$); tailoring, adding elements, lengthening, and cultural modifications ($k = 3$); and tailoring and cultural modifications.

Steps in adaptation

Each of the nine steps derived from the scoping review of evaluation frameworks [23] is represented in most of the adaptations (combining consulting with experts and stakeholders), with fewer reporting selecting the EBI ($k = 23$; 54.8%) and pilot testing ($k = 24$; 57.1%) (Table 3); 37 (88.1%) conducting a community assessment; 37 (88.1%) preparing new materials; 35 (83.3%) implementing the adapted intervention; 32 (76.2%) evaluating the adapted intervention; 31 (73.8%) determining needed changes based on action


step assessments; 31 (73.8%) training staff members; and 30 (71.4%) consulting stakeholders or experts before adapting the materials (Fig. 3). Overall, the average number of steps was 6.7 (range 3–9, mode = 7). Of the 37 authors who reported conducting community assessments, 21 (56.8%) held focus groups with community members, 12 (32.4%) conducted interviews with key informants and stakeholders, five (13.5%) formed and consulted with community advisory boards or steering committees, and two (5.4%) administered a survey to get community feedback. Ten of these (27.0%) used a combination of methods to collect community input and assess need.

Use of adaptation frameworks

The authors of less than half of the reports named a pre-existing adaptation framework as guiding the adaptation process ($k = 15$; 35.7%) (Table 2). Most frameworks were mentioned once; the Ecological Validity Model, Map of the Adaptation Process, and Cultural Adaptation Framework were referenced twice.

Intervention outcomes

Of the 36 reports that included an evaluation, most authors reported measuring program acceptability ($k = 28$; 66.7%), fidelity ($k = 22$; 52.4%), and feasibility ($k = 22$; 52.4%) (Table 3). With respect to implementation outcomes, most authors reported evaluating program acceptability ($k = 28$; 66.7%), fidelity ($k = 22$; 52.4%), and feasibility ($k = 22$; 52.4%). Several studies also reported assessing the adoption/implementation ($k = 20$; 47.6%) and sustainability of the program ($k = 11$; 26.2%). Numerous authors also reported measuring behavioral and






Fig. 3 Steps taken in the adaptation process across studies

program outcomes. The majority reported measuring behavioral outcomes ($k = 30$; 71.4%), while a smaller number measured changes in practice ($k = 9$; 21.4%), knowledge ($k = 7$; 16.7%), self-efficacy ($k = 5$; 11.9%), or environment ($k = 4$; 9.5%). Only a few studies included assessments of changes in well-being ($k = 3$; 7.1%), attitudes ($k = 3$; 7.1%), skills ($k = 3$; 7.1%), or communication ($n = 2$; 4.8%). Lastly, 11 (26.2%) of the evaluations included satisfaction with the adapted intervention as an outcome (Table 4).

Presentation of adapted elements

The authors used a variety of formats to present their adaptation processes. All 42 adaptations were described in the article's narrative, while others also used tables and figures to present certain elements. Seventeen (40.5%) included a table of the adaptations or modifications made. Three adaptations (7.1%) illustrated the adaptation process with a figure, and two (4.8%) included a side-by-side comparison of the adapted and original EBIs.

Discussion

This study presents findings based on a systematic review of published reports of adaptations of 42 EBIs. We present a systematic characterization of reasons for adapting EBIs, types of modifications made, steps taken during adaptation, reference to existing adaptation frameworks, and the constructs measured in evaluations of the adapted EBIs. In our review, the most common reasons for adaptation were to be relevant to a particular culture or new population, and to implement a program in a new setting. A previous study by Moore also found cultural adaptation to be a common reason for adaptations among evidence-based grantees, although less frequently (43% compared to our 64%) [15]. Higher frequency reasons in Moore's study were related to

resource constraints or logistics: lack of time (80%), limited resources (72%), difficulty retaining participants (71%), and resistance from implementers (64%) [15].

Among our included reports, all adaptation teams, or individuals involved in the research or adaptation, conducted content modifications, most commonly tailoring, adding or removing elements, and shortening. In their review of 32 published descriptions of interventions implemented in routine care or community settings, Stirman and colleagues also found the same four content modifications most frequently reported [1]. Consistent with the Stirman review, we found that context modifications were the next most frequently mentioned type of adaptation for either the program population or setting. Stirman, however, also found that format changes were frequently described [1]. In our review, delivery modifications were described in the majority of the studies, with training and evaluation modifications much less common. It is unclear whether these did not occur or were less often reported. Moore's review found slightly different frequencies of modifications, with more reports of changes related to logistics such as changes in delivery and dose, and much less frequent content changes [15].

Like Krivitsky, we also found that 29% reported removing elements [34]. This type of adaptation should be explored more because of its implications for reducing the fidelity to program core elements and potentially reducing the EBI's effects [35]. Additionally, four studies explicitly described changing the core elements of the original EBI. This is an area of concern because the integrity of the original program could have been jeopardized. More research is needed to understand why the elements were deleted and if the program implementers (i.e., researchers, community planners) consulted others before undertaking this change. The low reporting of changes to core elements may be because it is difficult

Table 4 Characteristics of the adaptations ($k = 42$)

First author, year	Adaptation type ¹					Specific modifications			Adaptation steps ²										Evaluation outcomes ($k = 36$)	Modification/adaptation example
	1	2	3	4	5	Content	Context	Delivery	1	2	3	4	5	6	7	8	9	10		
Reijneveld, 2003 [53]	x	x	x	x	x	Tailoring Adding elements Lengthening Substitution Cultural modification	Population	Deliverer Other	x	x		x	x	x	x	x	x	x	Acceptability Fidelity	Examples regarding safety excluded cycling because few Turkish immigrants cycle
Komro, 2004 [54]	x	x	x			Tailoring Adding elements Cultural modification	Setting Population	Mode/ medium Deliverer	x	x	x	x	x	x	x	x	x		Acceptability Fidelity	Audiotape vignettes re-taped with African American and Hispanic actors
Sarkisian, 2005 [55]	x			x		Tailoring Adding elements Lengthening Cultural modification	Setting Population	–	x		x		x						Acceptability	Expanded focus to more explicitly include family members
Tsey, 2005 [56]	x	x	x			Tailoring Shortening Loosening structure	Setting Population	Mode/ medium Deliverer	x								x	x	Acceptability Adoption Sustainability Individual satisfaction	Students interviewed their role models, explaining why they looked up to that person
Villarruel, 2005 [58]	x	x		x	x	Tailoring Adding elements Cultural modification	Population	–	x	x	x	x	x	x	x	x	x	x	Acceptability Individual satisfaction	Presented the view of machismo that incorporated the values of caring for and protecting others, so condom use could be presented as consistent with machismo
Belansky, 2006 [59]	x	x		x	x	Tailoring Adding elements Shortening Cultural modification	Setting Population	–	x		x	x	x	x	x	x	x	x	–	Lessons simplified so that they could be completed during the 1-h classroom period
Hitt, 2006 [60]	x	x		x		Tailoring Loosening structure	Setting Population	–	x	x	x	x	x	x	x		x	x	Acceptability Adoption Fidelity Feasibility Sustainability Individual satisfaction	Intervention protocols and supporting materials (tools) were tailored for local circumstances
Somerville, 2006 [62]	x	x		x	x	Tailoring Adding elements Substitution Integrating other approach Cultural modification	Setting Population	–	x	x	x	x	x	x	x	x	x	x	Acceptability Fidelity Other Individual satisfaction	A variety of successful Latino-focused HIV prevention training programs were integrated into the adapted intervention
NIMH Collaborative HIV/STD Prevention Trial Group, 2007 [38]	x	x		x	x	Tailoring Cultural modification	Other	–	x	x	x	x	x	x	x	x	x	x	Adoption Fidelity Feasibility Sustainability	Specific messages used in training were based on findings that emerged from the ethnography with each site's populations
Tsarouk, 2007 [63]	x	x				Tailoring Removing elements Shortening Substitution Cultural modification	Other	–	x		x		x	x	x		x		Acceptability Feasibility Individual satisfaction	Some of the support behaviors, such as applauding in response to a group member's participation, were removed because teens said that it is not a natural expression of support in this informal situation

Table 4 Characteristics of the adaptations ($k = 42$) (Continued)

First author, year	Adaptation type ¹					Specific modifications			Adaptation steps ²										Evaluation outcomes ($k = 36$)	Modification/adaptation example
	1	2	3	4	5	Content	Context	Delivery	1	2	3	4	5	6	7	8	9	10		
Beattie, 2008 [64]	x	x			x	Tailoring Removing elements Shortening Integrating other approach Cultural modification	Setting Population	–	x		x	x	x			x	x		Acceptability Adoption Sustainability	Some sites used a swim camp model, with several days of training provided often on two or three occasions and typically at a central point for families traveling long distances
Cornelius, 2008 [28]		x	x			Tailoring Adding elements Removing elements Lengthening Substitution	Setting Population	–	x	x				x					–	Used videos that included information about HIV in older women
Gitlin, 2008 [30]	x	x	x			Tailoring Adding elements Cultural modification	Setting Population	–	x		x	x		x		x	x		Acceptability Adoption Fidelity Feasibility Sustainability Other Individual satisfaction	Introduction of moment of silence at the beginning of each session to recognize spiritual practices and their importance to participants
Lerdboon, 2008 [67]		x	x			Tailoring Adding elements Integrating other approach Cultural modification	Setting	–	x	x	x	x	x	x	x	x	x	x	Acceptability Adoption Feasibility	Gender-specific components were integrated throughout the curriculum through a story line about an adolescent boy and girl growing up in a fictional Vietnamese family, as well as gender-specific scenarios, activities and messages
Steiker, 2008 [29]		x	x			Tailoring Adding elements Cultural modification	Setting Population	–	x		x		x	x	x	x	x	x	Acceptability Adoption Fidelity	Created four new videos, one for each prevention strategy: refuse, explain, avoid and leave
Burgio, 2009 [68]		x	x	x	x	Tailoring Removing elements Shortening	Setting Population	Mode/ medium Deliverer	x		x	x	x	x		x	x		Acceptability Adoption Fidelity Feasibility Individual satisfaction	Reduced number of home visits and shortened time span of the intervention
Fiscian, 2009 [69]		x	x	x	x	Tailoring Adding elements Removing elements Lengthening Reorder elements Integrating other approach Cultural modification	Setting Population	Mode/ medium	x		x	x	x	x		x	x		Acceptability Fidelity Individual satisfaction	Modified role-play stories to use African names and settings and simplified scripts to a sixth-grade reading level
Mueller, 2009 [31]		x	x			Adding elements Lengthening Substitution Loosening structure	Setting	–	x		x	x	x	x		x			Acceptability Feasibility Sustainability Individual satisfaction	Adapted from community agency or after-school programs to be integrated into existing school curriculum
Pekmez, 2009 [70]		x	x		x	Tailoring Cultural modification	Population	–	x		x			x	x	x	x		Acceptability Feasibility Individual satisfaction	Intervention materials and research measures were translated into Spanish through an iterative process involving both translation and back-translation
Stevens, 2009 [71]		x	x	x	x	Tailoring Adding elements	–	Deliverer		x	x	x	x	x		x	x		Fidelity	Support teams for caregivers were created
DePue, 2010 [72]		x	x	x	x	Tailoring Adding elements Loosening structure Cultural modification	Setting Population	Mode/ medium Deliverer	x	x	x	x	x	x	x	x	x	x	Adoption Feasibility	Incorporated local cultural features in flipcharts, including quotes from focus groups, culturally relevant examples of healthy behaviors, local sources of stress, and effective local coping strategies

Table 4 Characteristics of the adaptations ($k = 42$) (Continued)

First author, year	Adaptation type ¹					Specific modifications			Adaptation steps ²										Evaluation outcomes ($k = 36$)	Modification/adaptation example
	1	2	3	4	5				1	2	3	4	5	6	7	8	9	10		
Domenech Rodriguez, 2011 [73]	x	x	x			Tailoring Adding elements Loosening structure Cultural modification	Setting Population	Deliverer	x	x	x	x	x	x	x	x	x	x	Acceptability Fidelity Feasibility Individual satisfaction	Sayings, or <i>dichos</i> , were incorporated generously into treatment manual as parents used them during the parent training sessions
Poulsen, 2010 [39] Vandenhout, 2010 [75]	x	x	x			Tailoring Adding elements Substitution Cultural modification	Setting Population Other	Mode/ medium Deliverer	x	x	x	x	x	x	x	x	x	x	Acceptability Adoption Fidelity Sustainability Individual satisfaction	Owing to low literacy rates among local adults, drawings were used to illustrate messages that were originally conveyed through text on posters and handouts
Sadler, 2010 [76]	x					Tailoring Adding elements Shortening Reorder elements Cultural modification	Population	Deliverer	x						x	x			–	PowerPoint voice over changed to be in the first person instead of third to inspire comradery and motivation for women battling cancer together through clinical trials
Rotheram-Borus, 2011 [77]	x	x	x			Tailoring Adding elements Removing elements Shortening Substitution Integrating intervention Repeating elements Cultural modification	Setting Population	Mode/ medium Deliverer	x						x	x	x	x	Other	The intervention content and framing was adapted to resonate with Buddhist values and idioms around “sound body and sound mind”, as well as Thai values around the importance of family and community in health and well-being
Cardona, 2009 [78]	x	x	x			Tailoring Adding elements Loosening structure Cultural modification	Setting Population	Other	x	x	x	x	x	x			x	x	Acceptability Adoption Fidelity Feasibility Individual satisfaction	Substituted a booster session with a session on “Parenting between two cultures” to add relevance to Latino immigrant families
Feinberg, 2012 [79]	x	x	x	x		Tailoring Removing elements Integrating intervention Cultural modification	Setting Population	Mode/ medium Deliverer Other	x	x	x	x	x	x	x	x	x	x	Acceptability Fidelity Feasibility Individual satisfaction	Reframed the focus of the intervention from prevention of depression to learning new skills to deal with everyday stress, with an emphasis on parenting
Parker, 2012 [81] Chen, 2013 [82]	x	x	x			Tailoring Adding elements Removing elements Lengthening Reorder elements Cultural modification	Population	Mode/ medium	x	x	x				x	x			Acceptability Adoption Fidelity Feasibility	Created “action plan for sustainability” to link participants with exercise/disease self-management programs in neighborhood
Reid, 2012 [83]	x	x				Tailoring Adding elements Departing Cultural modification	Setting Population	Other	x		x	x	x	x	x	x	x		Acceptability Adoption Fidelity Feasibility Individual satisfaction	Sociocultural norms, values, beliefs, and myths were applied to role-play scenarios and exercises
Rosati, 2012 [84]	x	x				Tailoring Adding elements Loosening structure Cultural modification	Setting	Mode/ medium	x	x		x			x		x	x	Acceptability Adoption Fidelity Feasibility Individual satisfaction	Added a unit targeting adolescent dating and sexual behavior after conducting focus groups with Thai parents
Tomioka, 2012 [85]	x	x		x	x	Tailoring Adding elements Lengthening Cultural modification	Population	–	x	x	x	x			x		x	x	Acceptability Adoption Fidelity Feasibility Sustainability Individual satisfaction	Added opening session with a prayer, a 6-month reunion, and provided certificate of completion

Table 4 Characteristics of the adaptations ($k = 42$) (Continued)

First author, year	Adaptation type ¹					Specific modifications			Adaptation steps ²										Evaluation outcomes (k = 36)	Modification/adaptation example
	1	2	3	4	5				1	2	3	4	5	6	7	8	9	10		
Danielson, 2013 [33]	x	x	x		x	Tailoring Shortening Loosening structure Departing	Setting Population	Mode/ medium					x	x		x	x		Adoption Fidelity Feasibility Other Individual satisfaction	Used a web-based delivery platform instead of small group sessions with 10–12 girls
Fasula, 2013 [86]	x	x	x	x		Tailoring Adding elements Shortening Lengthening Loosening structure Cultural modification	Setting Population	Deliverer	x	x	x	x	x	x	x				–	Several intervention elements were added to increase participants' risk awareness, knowledge, and skills related to substance use, including a group discussion about the pros and cons of substance use, how drugs/alcohol contribute to sexual risk, and strategies for avoiding risk
Parker, 2013a [88] Parker, 2013b [89]	x	x	x			Tailoring Adding elements Removing elements Shortening Integrating other approach Cultural modification	Setting Population	Mode/ medium	x	x	x	x		x	x	x	x		Acceptability Feasibility	Changed delivery from individual to group so there was peer reinforcement content
Wainer, 2013 [90]	x	x	x	x	x	Shortening Loosening structure	Setting	Mode/ medium					x		x	x		x	Acceptability Fidelity Feasibility	Therapists completed the online training program on computers in their homes or in the research lab
Williams, 2013 [91]	x	x	x	x		Tailoring Adding elements Cultural modification	Setting Population	Deliverer	x	x	x	x	x			x	x		Adoption	The culturally adapted intervention took a more deliberate and structured approach to including the family in discussion and planning
Baydala, 2014 [92]	x	x			x	Tailoring Adding elements Lengthening Cultural modification	Population	–	x	x	x	x	x	x	x	x	x	x	Acceptability Adoption Fidelity Feasibility Sustainability	Elders suggested inclusion of lessons that embraced Aboriginal spirituality, such as an activity on healing the worried mind where students were encouraged to take their worried mind to <i>Waka</i> (God/Creator) and engage in <i>wacigebi</i> (prayer)
Broning, 2014 [93]	x	x				Tailoring Cultural modification	Population	–					x			x	x	x	–	Intervention was translated and adapted to German culture, taking into account family-based interventions are especially culture-sensitive regarding role-model behavior, values and norms
Cariou, 2014 [94]	x	x		x	x	Tailoring Adding elements Removing elements	Setting Population	–	x	x	x	x	x	x		x	x	x	Adoption Sustainability Other	Eliminated optional poolside activities and retained the few that were feasible based on available resources
Reback, 2014 [96]	x	x	x	x	x	Tailoring Adding elements Removing elements Shortening Reorder elements Cultural modification	Setting	Deliverer	x		x	x	x	x		x	x		Adoption Feasibility Sustainability	Gay-specific cultural references were updated to maintain cultural relevancy (i.e., exchanging references to telephone dating lines with references to social networking web sites)

Table 4 Characteristics of the adaptations ($k = 42$) (Continued)

First author, year	Adaptation type ¹					Specific modifications			Adaptation steps ²										Evaluation outcomes ($k = 36$)	Modification/adaptation example
	1	2	3	4	5				1	2	3	4	5	6	7	8	9	10		
Riggs, 2014 [32]	x		x		x	Adding elements	–	Mode/medium	x	x			x	x	x	x	x		Acceptability	Masters-level research
						Shortening		Deliverer											Fidelity	interventionists delivered
																			Feasibility	treatment rather than
																				medical staff
Tu, 2014 [98]	x	x	x	x	x	Tailoring	Population	Deliverer				x		x		x	x	–		In-person education from
						Removing elements														health educator was deleted

¹Adaptation type: (1) content, (2) context, (3) delivery, (4) training, and (5) evaluation

²Adaptation steps: (1) community assessment, (2) selection, (3) determine level of change, (4) train staff, (5) consult stakeholders/experts, (6) prepare materials, (7) pilot, (8) implement, (9) evaluate, and (10) other

to identify what the core elements are in an EBI. They may include elements that are readily adapted such as delivery or content. However, unless the original developers of the program or health-related online clearing-houses or resources where they are housed clearly describe them, it is often difficult for planners to identify them. Therefore, considerations of fidelity are critical when making decisions about what to adapt [36]. In a systematic review by Gearing and colleagues of 24 meta-analyses and review articles focusing on fidelity over the past 30 years, the authors identified core components of fidelity including design, training, and monitoring of intervention receipt and suggested that greater attention is needed to document threats to fidelity that remain underreported [37]. While this is true for any implementation effort, it is even more important to consider when making and reporting adaptations. Of particular note in our findings, while many authors reported changes to the delivery of the EBI, including who delivered it, there were fewer who reported adapting training or monitoring of that delivery.

Although cultural modification is not part of Stirman's taxonomy of adaptation modifications, we found that almost 75% of the authors described their adaptation in this way. Because cultural adaptations would almost always require some adaptation related to population and context, it is likely that authors in the Stirman review reporting adaptations to content, context, and new populations were, at least in some cases, making cultural adaptations. More clarity in definitions of what is meant by each type of adaptations is needed.

Our review uses a new taxonomy of steps or tasks for adaptation derived from a scoping study of existing frameworks [23]. We looked for nine steps or tasks and found that two adaptations reported all of the steps [38, 39], with the mean number being seven. Thus, most adaptation teams completed the majority of the steps. Overall, we found that most reported community assessment, preparation of materials, implementation, evaluation, and engaging stakeholders/experts as part of a program planning process. The least common step was selection of the EBI. This may be because some program staff may have already

decided on the EBI a priori and did not undertake a process to review candidate EBIs and select one.

The Escoffery classification from a review of adaptation frameworks seems applicable to real-world adaptation and could be used by others as a taxonomy for describing adaptation steps [23]. However, there are details that may be nuanced that are important to understand for the field both in describing adaptation steps and for informing future adaptations of the same EBI. For example, some reports include information about which components of the intervention were pilot tested and what decisions were made based on assessment finding, who the stakeholders were, and how they were engaged. Additionally, specific details about who is involved in the adaptation process (stakeholders, target population, program deliverers, health promotion, and behavioral scientists) and who makes the final decisions on what changes to make are critical to document. This information could be very important in interpreting reasons for specific adaptations and informing subsequent ones for future EBI implementation.

Capacity building efforts can assist practitioners to document the process in more detail and be deliberative or proactive with adaptations. The Cancer Prevention and Control Research Network (CPCRN) has modules on program selection of EBIs and adaptation with tools that help practitioners to document the discussion and decisions related to those processes in their Putting Public Health Prevention into Practice training [40]. In addition, the new online decision support tool, IM ADAPT, walks public health professionals through a systematic process to create a logic model for the adapted EBI and a selection adaptation, implementation, and evaluation plan based on intervention mapping [41, 42].

Among those reports that included an evaluation, the most common outcomes were acceptability, fidelity, and feasibility. This is not surprising since acceptability and feasibility of an intervention is often associated with program adoption [27]. Only one third reported the use of an adaptation framework to inform their process. This number is surprisingly low. Adaptation frameworks would provide guidance and rationale for this process

and should be used. Many frameworks exist, but perhaps program planners may not be aware of them [34] or may not know how to follow them without training or technical assistance. Due to the limited research on program adaptation, there also may not be the emphasis on adaptation models and frameworks. There needs to be increased dissemination and education on these frameworks to offer assistance with recommended steps in program adaptation.

Implications

Through a search of the published literature, this is the first systematic review of adapted evidence-based public health interventions internationally. Findings from the present study lead to important implications for the field of implementation science. First, many of the reasons for adaptation focus on either a change of population or setting, while the most common modifications were related to content, context, and delivery. Program developers of EBIs could anticipate program adaptation, instead of only adoption, and provide technical guidance in making modifications in their implementation protocols (or facilitator's manual) or program website. Recognizing that it is likely that successful programs will be adapted, program developers should also provide guidance about the theory and mechanisms of change that were used in the intervention and where possible design flexibility to match various contexts and populations [43]. They also can serve as expert consultants to help in the adaptation process as recommended as part of adaptation steps in adaptation frameworks [11, 16, 44] or our scoping study [23]. In addition, they could support a community of their EBI adopters by making adapted versions available or offering contacts for practitioners implementing the same program. Due to the low reporting of use of adaptation frameworks, the frameworks could be more widely disseminated to inform future adaptation efforts.

Research on best methods to document program adaptation is warranted to better understand whether it is best to describe and code adaptations based on document reviews of adaptation plans, published articles or reports, interviews of the adaptation team, or all of the above. Each of these methods has limitations, but implementing them all may not be practical for research studies. Finally, we found a variety of styles in reporting the reasons, modifications made, and process of adaptation. Standardization of reporting elements on program adaptation would guide professionals in describing their changes to EBIs and advance the field. Through this process of better reporting on adaptations, practitioners and program planners can better understand the reasons for adaptation, the adaptation process, and results to inform their own practice. Currently, TREND and Standards for Reporting Implementation Studies (StaRI) statement only ask researchers

to report on adaptation in general or adaptation results [45, 46]. Other critical elements of adaptations that we have identified (i.e., reasons, types of adaptations, steps taken) are not mentioned or delineated. There is a growing body of literature of adaptation taxonomies that could be recommended for some of these elements, including types of modifications [1, 47], reasons, timing and valence [15, 47], frameworks employed, and steps taken [23]. Creation of detailed reporting standards for program adaptation will result in commonalities for describing adaptations in the published and gray literature and will advance the field of implementation science in terms of producing adaptation data for further analysis.

Future research could explore planned versus unplanned adaptations and patterns of program modifications and the reasons for that happening. We present some early findings of patterns of modifications made to public health EBIs, but there is scarce understanding of them. In addition, further evaluations of adapted interventions are required to determine if adapted versions are as effective as the original program or other adapted versions. In this study, over 60% of the adaptation reports were non-experimental (i.e., observational, pilot program, post-test only) and less than one third were experimental. It is important for the field to have more rigorous evaluations of adapted programs to understand their outcomes and if their effects are comparable, better, or worse than the original EBI. Some preliminary research suggests that adapted versions of interventions are not associated with worst outcomes [48]. The evaluations also could inform if different types or combinations of modifications (e.g., content and context) impact effectiveness as well. Researchers also should determine critical adaptation elements to record and standardize across studies. Finally, while there have been repositories of evidence-based programs for public health practice such as the National Registry for Effective Programs and Practices [49] and Research Tested Intervention Programs (RTIPs) [50], there is no clearinghouse for adapted programs for the field to understand the issues around external validity of EBIs. Chambers recommends the creation of an adaptome to catalog adapted programs and their results to share with the field to potentially address this gap [51].

Several limitations exist for this study. Although we searched for relevant articles of adapted EBIs, it is likely that some articles were overlooked based on our search strategy. For example, we did not review gray literature for adapted EBIs. In addition, we limited our studies to those that focus on public health interventions and excluded clinical interventions. Additionally, our data on adaptations made and outcomes were limited to the authors' description in the article and were not augmented with other data (e.g., surveys of authors). Although we

had two raters to increase the reliability of the data abstracted, some of the modifications may have been underreported if the authors fail to report on that aspect (i.e., training) or may not have fully implemented the program yet (i.e., evaluation). Finally, while our review included adapted interventions globally, we did not review articles in languages other than English. However, we were able to find 16 studies in international settings. A limitation of this study is that we did not confirm with the authors that all of the adaptations made were reported; for example, some could have not reported on unplanned adaptation since some were not yet implemented. It is helpful for planners to document all adaptation, both planned and unplanned [52], for other practitioners to learn from this process. Finally, this review is becoming dated, especially in an area with a much active research and reporting.

Conclusion

This review offers a practical examination of adaptation across multiple programs and program types that were implemented in community settings. It reports systematically on reasons for adaptation, types of modifications, and steps of adaptations for public health EBIs in public health practice. Adaptations are occurring in natural settings for a variety of reasons, and commonly, adaptations are made to intervention content or context. A few steps were used across adaptation teams in the process of adaptations, but the science of adaptation is still an emerging area of study in implementation science. More critical appraisal of intervention adaptations and their outcomes could assist with EBI transferability to increase the scale up and spread of EBI to increase population health impact.

Additional files

Additional file 1: Systematic review search terms. (DOCX 13 kb)

Additional file 2: Abstraction categories and definitions. (DOCX 19 kb)

Abbreviations

CPCRN: Cancer Prevention and Control Research Network; EBIs: Evidence-based interventions; MeSH: Medical Subject Headings

Acknowledgements

We are grateful to Danielle Reece, Shuting Liang, and Scott Decker for their contributions during the data abstraction phase of this study.

Funding

This research was supported in part by the Increasing Reach and Implementation of Evidence Based Programs for Cancer Control, National Cancer Institute grant (R01-CA163526) and the Cancer Prevention and Control Research Network (3 U48 DP005017-01S8). The content of this paper is solely the responsibility of the authors and does not necessarily represent the official views of the funding agency. No financial disclosures were reported by the authors of this paper.

Availability of data and materials

Please contact the authors for data requests.

Authors' contributions

All authors were involved in various stages of the study design. CE conceptualized the study, and MEF, MH, and PDM helped to design the study questions and abstraction tool. CE, EL, EB, and HU wrote the first draft. All abstracted the articles, assisted with the data interpretation, and commented on the subsequent drafts of the paper. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Rollins School of Public Health, Emory University, 1518 Clifton Road, Atlanta, GA 30322, USA. ²University of Texas School of Public Health, 7000 Fannin, Ste 2522, Houston, TX 77030, USA.

Received: 22 February 2018 Accepted: 10 September 2018

Published online: 26 September 2018

References

1. Stirman SW, Miller CJ, Toder K, Calloway A. Development of a framework and coding system for modifications and adaptations of evidence-based interventions. *Implement Sci*. 2013;8:1.
2. Durlak JA, DuPre EP. Implementation matters: a review of research on the influence of implementation on program outcomes and the factors affecting implementation. *Am J Community Psychol*. 2008;41:327–50.
3. Card JJ, Solomon J, Cunningham SD. How to adapt effective programs for use in new contexts. *Health Promot Pract*. 2011;12:25–35.
4. Bell A, Corfield M, Davies J, Richardson N. Collaborative transdisciplinary intervention in early years - putting theory into practice. *Child Care Health Dev*. 2010;36:142–8.
5. Rogers E. Diffusion of innovations. New York: Simon & Schuster; 1995.
6. Solomon J, Card JJ, Malow RM. Adapting efficacious interventions: advancing translational research in HIV prevention. *Eval Health Prof*. 2006;29:162–94.
7. Lee SJ, Altschul I, Mowbray CT. Using planned adaptation to implement evidence-based programs with new populations. *Am J Community Psychol*. 2008;41:290–303.
8. Chen EK, Reid M, Parker SJ, Pillemer K. Tailoring evidence-based interventions for new populations: a method for program adaptation through community engagement. *Eval Health Prof*. 2012;36(1):73–92.
9. O'Connor C, Small, SA. Program fidelity and adaptation: Meeting local needs without compromising program effectiveness. https://fyi.uwex.edu/whatworkswisconsin/files/2014/04/whatworks_04.pdf. Accessed 17 Sept 2018.
10. Roller LA, Fuller TR, Firpo-Triplett R, Lesesne CA, Moore C, Leeks KD. Adaptation guidance for evidence-based teen pregnancy and STI/HIV prevention curricula: from development to practice. *Am J Sex Educ*. 2014;9:135–54.
11. Backer T. Finding the balance: program fidelity and adaptation in substance abuse prevention: a state-of-the-art review. Rockville: Center for Substance Abuse Prevention. Substance Abuse and Mental Health Services Administration; 2002.
12. Bartholomew Eldredge L, Highfield L, Hartman M, Mullen P, Leerlooijer J, Fernandez M. Using intervention mapping to adapt evidence-based interventions. In: Bartholomew Eldredge L, Markham C, Ruiter R, Fernandez M, Kok G, Parcel G, editors. Planning health promotion programs: an intervention mapping approach. San Francisco: John Wiley & Sons; 2016. p. 596–649.
13. Bartholomew L, Parcel G, Kok G, Gottlieb N. Planning health promotion programs: an intervention mapping approach. San Francisco: Jossey-Bass; 2006.
14. Smith E, Caldwell L. Adapting evidence-based programs to new contexts: what needs to be changed? *J Rural Health*. 2007;23(Suppl):37–41.
15. Moore JE, Bumbarger BK, Cooper BR. Examining adaptations of evidence-based programs in natural contexts. *J Prim Prev*. 2013;34:147–61.

16. McKleroy VS, Galbraith JS, Cummings B, Jones P, Harshbarger C, Collins C, Gelaude D, Carey JW. Adapting evidence-based behavioral interventions for new settings and target populations. *AIDS Educ Prev*. 2006;18:59–73.
17. Barrera M, Castro FG. A heuristic framework for the cultural adaptation of interventions. *Clin. Psychol. Sci. Pract.* 2006;13:311–6.
18. Carvalho ML, Honeycutt S, Escoffery C, Glanz K, Sabbs D, Kegler MC. Balancing fidelity and adaptation: implementing evidence-based chronic disease prevention programs. *J Public Health Manag Pract*. 2013;19(4):348–56.
19. Cooper BR, Shrestha G, Hyman L, Hill L. Adaptations in a community-based family intervention: replication of two coding schemes. *J Prim Prev*. 2016;37:33–52.
20. Cohen DJ, Crabtree BF, Etz RS, Balasubramanian BA, Donahue KE, Leviton LC, Clark EC, Isaacson NF, Stange KC, Green LW. Fidelity versus flexibility: translating evidence-based research into practice. *Am J Prev Med*. 2008;35:S381–9.
21. Miller-Day M, Pettigrew J, Hecht ML, Shin Y, Graham J, Krieger J. How prevention curricula are taught under real-world conditions: types of and reasons for teacher curriculum adaptations. *Health Educ*. 2013;113:324–44.
22. Kumpfer KL, Pinyuchon M, Teixeira de Melo A, Whiteside HO. Cultural adaptation process for international dissemination of the strengthening families program. *Eval Health Prof*. 2008;31:226–39.
23. Escoffery C, Lebow-Skelley E, Udelson H, Boing EA, Wood R, Fernandez ME, Mullen PD. A scoping study of frameworks for adapting public health evidence-based interventions. *Transl Behav Med*. 2018. in press.
24. Wandersman A, Duffy J, Flaspohler P, Noonan R, Lubell K, Stillman L, Blachman M, Dunville R, Saul J. Bridging the gap between prevention research and practice: the interactive systems framework for dissemination and implementation. *Am J Community Psychol*. 2008;41:171–81.
25. Higgins JPT, Green S. *Cochrane handbook for systematic reviews of interventions* version 5.1, vol. 0. Chichester: The Cochrane Collaboration; 2011.
26. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol*. 2009;62:1006–12.
27. Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, Griffey R, Hensley M. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Adm Policy Ment Health*. 2011;38:65–76.
28. Cornelius JB, Moneyham L, LeGrand S. Adaptation of an HIV prevention curriculum for use with older african american women. *J Assoc Nurses AIDS Care*. 2008;19:16–27.
29. Steiker LKH. Making drug and alcohol prevention relevant: adapting evidence-based curricula to unique adolescent cultures. *Fam Community Health*. 2008;31:S52–60.
30. Gitlin LN, Chernett NL, Harris LF, Palmer D, Hopkins P, Dennis MP. Translation of the chronic disease self-management program for older african americans in a senior setting. *Gerontologist*. 2008;48:698–705.
31. Mueller TE, Castaneda CA, Sainer S, Martinez D, Herbst JH, Wilkes AL, Villarruel AM. The implementation of a culturally based hiv sexual risk reduction program for latino youth in a Denver area high school. *AIDS Educ Prev*. 2009;21:164–70.
32. Riggs KR, Lozano P, Mohelnitzky A, Rudnick S, Richards J. An adaptation of family-based behavioral pediatric obesity treatment for a primary care setting: group health family wellness program pilot. *Perm J*. 2014;18:4–10.
33. Danielson CK, McCauley JL, Jones AM, Borkman AL, Miller S, Ruggiero KJ. Feasibility of delivering evidence-based HIV/STI prevention programming to a community sample of African American teen girls via the internet. *AIDS Educ Prev*. 2013;25:394–404.
34. Krivitsky LN, Parker SJ, Pal A, Meckler L, Shengelia R, Reid MC. A systematic review of health promotion and disease prevention program adaptations: how are programs adapted? In: Wethington E, Dunifon RE, editors. *APA Bronfenbrenner series on the ecology of human development. Research for the public good: applying the methods of translational research to improve human health and well-being*. Washington: American Psychological Association; 2012. p. 73–99.
35. Eke AN, Neumann MS, Wilkes AL, Jones PL. Preparing effective behavioral interventions to be used by prevention providers: the role of researchers during HIV prevention research trials. *AIDS Educ Prev*. 2006;18:44–58.
36. Bopp M, Saunders RP, Lattimore D. The tug-of-war: fidelity versus adaptation throughout the health promotion program life cycle. *J Prim Prev*. 2013;34:193–207.
37. Gearing RE, El-Bassel N, Ghesquiere A, Baldwin S, Gillies J, Ngeow E. Major ingredients of fidelity: a review and scientific guide to improving quality of intervention research implementation. *Clin Psychol Rev*. 2011;31:79–88.
38. NIMH Collaborative HIV/STD Prevention Trial Group. Methodological overview of a five-country community-level HIV/sexually transmitted disease prevention trial. *AIDS*. 2007;21:S3–S18.
39. Poulsen MN, Vandenhoude H, Wyckoff SC, Obong'o CO, Ochura J, Njika G, Otway NJ, Miller KS. Cultural adaptation of a U.S. evidence-based parenting intervention for rural Western Kenya: from parents matter! To families matter! *AIDS Educ Prev*. 2010;22:273–85.
40. Cancer Prevention and Control Research Network: Cancer Prevention and Control Research Network. 2018. <http://cpcrn.org/about/>. Accessed 5 Jan 2018.
41. Mullen PD, Bartholomew LK. Intervention Mapping step 6: planning for evaluation. In: Bartholomew LK, Parcel GS, Kok G, Gottlieb NH, Fernandez ME, editors. *Planning health promotion programs: an intervention mapping approach*. San Francisco: Jossey-Bass; 2011.
42. Bartholomew Eldredge LK, Highfield L, Hartman M, Mullen PD, Leerlooijer J, Fernandez M. Chp 10, Int Map Adapt: using intervention mapping to adapt evidence-based interventions. In: Bartholomew Eldredge LK, Markham C, RAC R, Fernandez ME, Kok G, Parcel GS, editors. *Planning health promotion programs: an intervention mapping approach*. San Francisco: Jossey-Bass; 2016.
43. Ham B, Parisi D, Stoolmiller M. Balancing fidelity with flexibility and fit: what do we really know about fidelity of implementation in schools? *Except Child*. 2013;79:181–93.
44. Wingood GM, DiClemente RJ. The ADAPT-ITT model: a novel method of adapting evidence-based HIV Interventions. *J Acquir Immune Defic Syndr*. 2008;47:S40–6.
45. Des Jarlais DC, Lyles C, Crepaz N, Group T. Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement. *Am J Public Health*. 2004;94:361–6.
46. Pinnock H, Barwick M, Carpenter CR, Eldridge S, Grandes G, Griffiths CJ, Rycroft-Malone J, Meissner P, Murray E, Patel A, et al. Standards for Reporting Implementation Studies (StaRI) statement. *BMJ*. 2017;6795:356.
47. Hill LG, Owens RW. Component analysis of adherence in a family intervention. *Health Educ*. 2013;113:264–80.
48. Stirman S, Gamarra J, Bartlett B, Calloway A, Gutner C. Empirical examinations of modifications and adaptations to evidence-based psychotherapies: methodologies, impact, and future directions. *Clin Psychol Sci Pract*. 2017;24(4):396–420.
49. Substance Abuse and Mental Health Services Administration: National Registry for Effective Programs and Practices. 2014. <https://www.samhsa.gov/nrepp>. Accessed 2 Jan 2018.
50. National Cancer Institute: Research Tested Intervention Programs (RTIPs). 2018. <https://rtips.cancer.gov/rtips/index.do>. Accessed 2 Jan 2018.
51. Chambers DA, Norton WE. The adaptome: advancing the science of intervention adaptation. *Am J Prev Med*. 2016;51:S124–31.
52. Galbraith JS, Stanton B, Boekeloo B, King W, Desmond S, Howard D, Black MM, Carey JW. Exploring implementation and fidelity of evidence-based behavioral interventions for HIV prevention: lessons learned from the focus on kids diffusion case study. *Health Educ Behav*. 2009;36:532–49.
53. Reijneveld SA, Westhoff MH, Hopman-Rock M. Promotion of health and physical activity improves the mental health of elderly immigrants: results of a group randomised controlled trial among Turkish immigrants in the Netherlands aged 45 and over. *J Epidemiol Community Health*. 2003;57:405–11.
54. Komro KA, Perry CL, Veblen-Mortenson S, Bosma LM, Dudovitz BS, Williams CL, Jones-Webb R, Toomey TL. The adaptation of Project Northland for urban youth. *J Pediatr Psychol*. 2004;29:457–66.
55. Sarkisian CA, Bruselas RJ, Steers WN, Davidson MB, Brown AF, Norris KC, Anderson RM, Mangione CM. Using focus groups of older African Americans and Latinos with diabetes to modify a self-care empowerment intervention. *Ethn Dis*. 2005;15:283–91.
56. Tsey K, Whiteside M, Daly S, Deemal A, Gibson T, Cadet-James Y, Wilson A, Santhanam R, Haswell-Elkins MR. Adapting the 'Family Wellbeing' empowerment program to the needs of remote Indigenous school children. *Aust N Z J Public Health*. 2005;29:112–6.
57. Tsey K, Deemal A, Whiteside M, Gibson T. It's an eye-opener for people to see you can change things: an evaluation of a 'family wellbeing' personal and community development pilot project in two Cape York Communities. St. Lucia: University of Queensland School of Population Health; 2003.
58. Villarruel AM, Jemmott LS, Jemmott JB. Designing a culturally based intervention to reduce HIV sexual risk for Latino adolescents. *J Assoc Nurses AIDS Care*. 2005;16:23–31.

59. Belansky ES, Romaniello C, Morin C, Uyeki T, Sawyer R, Scarbro S, et al. Adapting and implementing a long-term nutrition and physical activity curriculum to a rural, low-income, biethnic community. *J Nutr Educ Behav*. 2006;38:106–13.
60. Hitt JC, Robbins AS, Galbraith JS, Todd JD, Patel-Larson A, McFarlane JR, Spikes P, Carey JW. Adaptation and implementation of an evidence-based prevention counseling intervention in Texas. *AIDS Educ Prev*. 2006;18:108–18.
61. Kamb ML, Fishbein M, Douglas JM Jr, Rhodes F, Rogers J, Bolan G, et al. Efficacy of risk-reduction counseling to prevent human immunodeficiency virus and sexually transmitted diseases: a randomized controlled trial. Project RESPECT Study Group. *JAMA*. 1998;280:1161–7.
62. Somerville GG, Diaz S, Davis S, Coleman KD, Taveras S. Adapting the popular opinion leader intervention for Latino young migrant men who have sex with men. *AIDS Educ Prev*. 2006;18:137–48.
63. Tsarouk TT, Elaine A, Herting JR, Walsh E, Randell B. Culturally specific adaptation of a prevention intervention: an international collaborative research project. *Addict Behav*. 2007;32:1565–81.
64. Beattie N, Shaw P, Larson A. Water safety in the bush: strategies for addressing training needs in remote areas. *Rural Remote Health*. 2008;8:855.
65. DiClemente RJ, Wingood GM. A randomized controlled trial of an HIV sexual risk-reduction intervention for young African-American women. *JAMA*. 1995; 274:1271–6.
66. Lorig KR, Sobel DS, Stewart AL, Brown BW Jr, Bandura A, Ritter P, Gonzalez VM, Laurent DD, Holman HR. Evidence suggesting that a chronic disease self-management program can improve health status while reducing hospitalization: a randomized trial. *Med Care*. 1999;37:5–14.
67. Lerdboon P, Pham V, Green M, Riel R, Tho le H, Ha NT, Kaljee LM. Strategies for developing gender-specific HIV prevention for adolescents in Vietnam. *AIDS Educ Prev*. 2008;20:384–98.
68. Burgio LDC, Irene B, Schmid B, Wharton T, McCallum D, DeCoster J. Translating the REACH Caregiver Intervention for use by area agency on aging personnel: the REACH OUT Program. *Gerontologist*. 2009;49:103–16.
69. Fiscian VS, Obeng EK, Goldstein K, Shea JA, Turner BJ. Adapting a multifaceted U.S. HIV prevention education program for girls in Ghana. *AIDS Educ Prev*. 2009;21:67–79.
70. Pekmezi DW, Neighbors CJ, Lee CS, Gans KM, Bock BC, Morrow KM, Marquez B, Dunsiger S, Marcus BH. A culturally adapted physical activity intervention for Latinas: a randomized controlled trial. *Am J Prev Med*. 2009;37:495.
71. Stevens AB, Lancer K, Smith ER, Allen L, McGhee R. Engaging communities in evidence-based interventions for dementia caregivers. *Fam Community Health*. 2009;32:S83–92.
72. DePue JD, Rosen RK, Batts-Turner M, Bereolos N, House M, Held RF, Nu'usolia O, Tuitele J, Goldstein MG, McGarvey ST. Cultural translation of interventions: diabetes care in American Samoa. *Am J Public Health*. 2010; 100:2085–93.
73. Domenech Rodriguez MM, Baumann AA, Schwartz AL. Cultural adaptation of an evidence based intervention: from theory to practice in a Latino/a community context. *Am J Community Psychol*. 2011;47:170–86.
74. Forgatch MS, DeGarmo DS. Parenting through change: an effective prevention program for single mothers. *J Consult Clin Psychol*. 1999;67:711–24.
75. Vandenhoudt H, Miller KS, Ochura J, Wyckoff SC, Obong'o CO, Otwoma NJ, Pousen MN, Menten J, Marum E, Buvé A. Evaluation of a U.S. evidence-based parenting intervention in rural western Kenya: from parents matter! to families matter! *AIDS Educ Prev*. 2010;22:328–43.
76. Sadler GR, Gonzalez J, Mumman M, Cullen L, LaHousse SF, Malcarne V, Conde V, Riley N. Adapting a program to inform African American and Hispanic American women about cancer clinical trials. *J Cancer Educ*. 2010; 25:142–5.
77. Rotheram-Borus MJ, Swendeman D, Lee SJ, Li L, Amani B, Nartey M. Interventions for families affected by HIV. *Transl Behav Med*. 2011;1:313–26.
78. Cardona JP, Holtrop K, Cordova D Jr, Escobar-Chew AR, Horsford S, Tams L, et al. "Queremos aprender": Latino immigrants' call to integrate cultural adaptation with best practice knowledge in a parenting intervention. *Fam Process*. 2009;48:211–31.
79. Feinberg E, Stein R, Diaz-Linhart Y, Egbert L, Beardslee W, Hegel MT, Silverstein M. Adaptation of problem-solving treatment for prevention of depression among low-income, culturally diverse mothers. *Fam Community Health*. 2012;35:57–67.
80. Gath D, Catalan J. The treatment of emotional disorders in general practice: psychological methods versus medication. *J Psychosom Res*. 1986;30:381–6.
81. Parker SJ, Chen EK, Pillemer K, Filiberto D, Laureano E, Piper J, Schwartz-Leeper J, Robbins L, Reid MC. Participatory adaptation of an evidence-based, arthritis self-management program: making changes to improve program fit. *Fam Community Health*. 2012;35:236.
82. Chen EK, Reid MC, Parker SJ, Pillemer K. Tailoring evidence-based interventions for new populations: a method for program adaptation through community engagement. *Eval Health Prof*. 2013;36:73–92.
83. Reid SD, Malow RM, Rosenberg R. Alcohol, drugs, sexual behavior, and HIV in Trinidad and Tobago—the way forward. *J Int Assoc Provid AIDS Care*. 2012;11:66–82.
84. Rosati MJ, Cupp PK, Chookhare W, Miller BA, Byrnes HF, Fongkaew W, Vanderhoff J, Chamratrithirong A, Rhucharoenpornpanich O, Atwood KA. Successful implementation of Thai family matters: strategies and implications. *Health Promot Pract*. 2012;13:555–63.
85. Tomioka M, Braun KL, Compton M, Tanoue L. Adapting Stanford's Chronic Disease Self-Management Program to Hawaii's multicultural population. *Gerontologist*. 2012;52(1):121–32.
86. Fasula AM, Fogel CI, Gelaude D, Carry M, Gaiter J, Parker S. Project power: adapting an evidence-based HIV/STI prevention intervention for incarcerated women. *AIDS Educ Prev*. 2013;25:203–15.
87. Shain RN, Piper JM, Newton ER, Perdue ST, Ramos R, Champion JD, Guerra FA. A randomized, controlled trial of a behavioral intervention to prevent sexually transmitted disease among minority women. *N Engl J Med*. 1999; 340:93–100.
88. Parker L, Maman S, Pettifor A, Chalachala JL, Edmonds A, Golin CE, Moracco K, Behets F. Adaptation of a U.S. evidence-based Positive Prevention intervention for youth living with HIV/AIDS in Kinshasa, Democratic Republic of the Congo. *Eval Program Plann*. 2013;36:124–35.
89. Parker L, Maman S, Pettifor A, Chalachala JL, Edmonds A, Golin CE, Moracco K, Behets F, Sympa Study T. Feasibility analysis of an evidence-based positive prevention intervention for youth living with HIV/AIDS in Kinshasa, Democratic Republic of the Congo. *AIDS Educ Prev*. 2013;25:135–50.
90. Wainer A, Ingersoll B. Disseminating ASD interventions. A pilot study of a distance learning program for parents and professionals. *J Autism Dev Disord*. 2013;43:11–24.
91. Williams AB, Honghong W, Burgess J, Xianhong L, Danvers K. Cultural adaptation of an evidence-based nursing intervention to improve medication adherence among people living with HIV/AIDS (PLWHA) in China. *Int J Nurs Stud*. 2013;50:487–94.
92. Baydala L, Fletcher F, Worrell S, Kajner T, Letendre S, Letendre L, Rasmussen C. Partnership, knowledge translation, and substance abuse prevention with a first nations community. *Prog Community Health Partnersh*. 2014;8:145–55.
93. Bröning S, Sack PM, Thomsen M, Stolle M, Wendell A, Stappenbeck J, Thomasius R. Implementing and evaluating the German adaptation of the "Strengthening Families Program 10 – 14" - a randomized-controlled multicentre study. *BMC Public Health*. 2014;14:83.
94. Cariou C, Gonzales M, Krebill H. Adapting and implementing an evidence-based sun-safety education program in rural Idaho, 2012. *Prev Chronic Dis*. 2014;11:E77.
95. Glanz K, Geller AC, Shigaki D, Maddock JE, Isneec MR. A randomized trial of skin cancer prevention in aquatic settings: the Pool Cool program. *Health Psychol*. 2002;21:579–87.
96. Reback CJ, Veniegas R, Shoptaw S. Getting off: development of a model program for gay and bisexual male methamphetamine users. *J Homosex*. 2014;61:540–53.
97. Saelens BE, Lozano P, Scholz K. A randomized clinical trial comparing delivery of behavioral pediatric obesity treatment using standard and enhanced motivational approaches. *J Pediatr Psychol*. 2013;38:954–64.
98. Tu SP, Chun A, Yasui Y, Kuniyuki A, Yip MP, Taylor V, Bastani R. Adaptation of an evidence-based intervention to promote colorectal cancer screening: a quasi-experimental study. *Implement Sci*. 2014;9:85.
99. Tu SP, Taylor V, Yasui Y, Chun A, Yip MP, Acorda E, et al. Promoting culturally appropriate colorectal cancer screening through a health educator. *Cancer*. 2006;107:959–66.